arXiv:1202.1237v3 [math.NT] 12 Feb 2012

JACOBSTHAL SUMS, LEGENDRE POLYNOMIALS AND BINARY QUADRATIC FORMS

Zhi-Hong Sun

School of Mathematical Sciences, Huaiyin Normal University,
Huaian, Jiangsu 223001, PR China
Email: zhihongsun@yahoo.com
Homepage: http://www.hytc.edu.cn/xsjl/szh

Abstract. Let $p>3$ be a prime and $m,n\in\mathbb{Z}$ with $p\nmid mn$. Built on the work of Morton, in the paper we prove the uniform congruence:

$$\sum_{x=0}^{p-1}\Big(\frac{x^3+mx+n}{p}\Big)$$
$$\equiv\begin{cases}-(-3m)^{\frac{p-1}{4}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{5}{12}}{k}\big(\frac{4m^3+27n^2}{4m^3}\big)^k \pmod p & \text{if } 4\mid p-1,\\ \frac{2m}{9n}\big(\frac{-3m}{p}\big)(-3m)^{\frac{p+1}{4}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{5}{12}}{k}\big(\frac{4m^3+27n^2}{4m^3}\big)^k \pmod p & \text{if } 4\mid p-3,\end{cases}$$

where $(\frac{a}{p})$ is the Legendre symbol. We also establish many congruences for $x \pmod p$, where x is given by $p=x^2+dy^2$ or $4p=x^2+dy^2$, and pose some conjectures on supercongruences modulo p^2 concerning binary quadratic forms.

MSC: Primary 11A07, Secondary 33C45, 11E25, 11A15, 05A10
Keywords: Congruence; Jacobsthal sum; Legendre polynomial; binary quadratic form

1. Introduction.

Let $\{P_n(x)\}$ be the Legendre polynomials given by

$$P_0(x)=1,\ P_1(x)=x,\ (n+1)P_{n+1}(x)=(2n+1)xP_n(x)-nP_{n-1}(x)\ (n\ge 1).$$

It is well known that (see [G, (3.132)-(3.133)])

$$(1.1)\qquad P_n(x)=\frac{1}{2^n}\sum_{k=0}^{[n/2]}\binom{n}{k}(-1)^k\binom{2n-2k}{n}x^{n-2k}=\frac{1}{2^n\cdot n!}\cdot\frac{d^n}{dx^n}(x^2-1)^n,$$

The author was supported by the Natural Sciences Foundation of China (grant no. 10971078).

where $[a]$ is the greatest integer not exceeding a. From (1.1) we see that

$$(1.2)\qquad P_n(-x)=(-1)^nP_n(x),\quad P_{2m+1}(0)=0\quad\text{and}\quad P_{2m}(0)=\frac{(-1)^m}{2^{2m}}\binom{2m}{m}.$$

We also have the following formula due to Murphy ([G, (3.135)]):

$$(1.3)\qquad P_n(x)=\sum_{k=0}^{n}\binom{n}{k}\binom{n+k}{k}\Big(\frac{x-1}{2}\Big)^k=\sum_{k=0}^{n}\binom{2k}{k}\binom{n+k}{2k}\Big(\frac{x-1}{2}\Big)^k.$$

We remark that $\binom{n}{k}\binom{n+k}{k}=\binom{2k}{k}\binom{n+k}{2k}$.

Let $\mathbb{Z}$ be the set of integers. For a prime p let $\mathbb{Z}_p$ be the set of rational numbers whose denominator is coprime to p. Based on the work of Morton ([BM],[M]), in [S3-S6] the author proved that for a prime $p>3$ and $t\in\mathbb{Z}_p$,

(1.4)
$$P_{\frac{p-1}{2}}(t)\equiv-\Big(\frac{-6}{p}\Big)\sum_{x=0}^{p-1}\Big(\frac{x^3-3(t^2+3)x+2t(t^2-9)}{p}\Big)\pmod p,$$

(1.5)
$$P_{[\frac p4]}(t)\equiv-\Big(\frac{6}{p}\Big)\sum_{x=0}^{p-1}\Big(\frac{x^3-\frac{3(3t+5)}{2}x+9t+7}{p}\Big)\pmod p,$$

(1.6)
$$P_{[\frac p3]}(t)\equiv-\Big(\frac{p}{3}\Big)\sum_{x=0}^{p-1}\Big(\frac{x^3+3(4t-5)x+2(2t^2-14t+11)}{p}\Big)\pmod p,$$

(1.7)
$$P_{[\frac p6]}(t)\equiv-\Big(\frac{3}{p}\Big)\sum_{x=0}^{p-1}\Big(\frac{x^3-3x+2t}{p}\Big)\pmod p,$$

where $\left(\frac{a}{p}\right)$ is the Legendre symbol.

For a prime $p>3$ let F_p be the field of p elements. For $m,n\in F_p$ we call the curve $E_p:\ y^2=x^3+mx+n$ an elliptic curve provided that the discriminant $4m^3+27n^2\neq0$ in F_p. Let $\#E_p(y^2=x^3+mx+n)$ be the number of points (including infinity) on the curve $E_p:\ y^2=x^3+mx+n$ over F_p. It is easily seen that (see for example [S1])

$$(1.8)\qquad \#E_p(y^2=x^3+mx+n)=p+1+\sum_{k=0}^{p-1}\Big(\frac{x^3+mx+n}{p}\Big).$$

The famous inequality due to Hasse states that

$$\Big|\sum_{k=0}^{p-1}\Big(\frac{x^3+mx+n}{p}\Big)\Big|\le2\sqrt p.$$

It is important and difficult to evaluate the general Jacobsthal sum $\sum_{k=0}^{p-1}(\frac{x^3+mx+n}{p})$. So far, the sum is only determined by using Deuring's theorem when the corresponding elliptic curve has complex multiplication (see [LM],[I]). In 2006, Morton[M] established a congruence for $\sum_{k=0}^{p-1}(\frac{x^3+mx+n}{p}) \pmod p$ by using Jacobi polynomials.

Let $p>3$ be a prime and $m,n\in\mathbb{Z}_p$ with $mn\not\equiv 0 \pmod p$. Built on the work of Morton, in the paper we prove the uniform congruence:

(1.9)

$$\begin{aligned}&\sum_{x=0}^{p-1}\Big(\frac{x^3+mx+n}{p}\Big)\\ &\equiv\begin{cases}-(-3m)^{\frac{p-1}{4}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{5}{12}}{k}(\frac{4m^3+27n^2}{4m^3})^k \pmod p & \text{if } 4\mid p-1,\\ \frac{2m}{9n}(\frac{-3m}{p})(-3m)^{\frac{p+1}{4}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{5}{12}}{k}(\frac{4m^3+27n^2}{4m^3})^k \pmod p & \text{if } 4\mid p-3.\end{cases}\\ &\equiv\begin{cases}(-1)^{\frac{p+1}{2}}(\frac{n}{2})^{\frac{p-1}{6}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{7}{12}}{k}(\frac{4m^3+27n^2}{27n^2})^k \pmod p & \text{if } 3\mid p-1,\\ (-1)^{\frac{p+1}{2}}\frac{3}{m}(\frac{2}{n})^{\frac{p-5}{6}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{7}{12}}{k}(\frac{4m^3+27n^2}{27n^2})^k \pmod p & \text{if } 3\mid p-2.\end{cases}\end{aligned}$$

We note that the right sums in (1.9) have the good form $\sum_{k=0}^{p-1}f(k)$.

For positive integers a,b and n, if $n=ax^2+by^2$ for some $x,y\in\mathbb{Z}$, we briefly say that $n=ax^2+by^2$. Let p be a prime of the form $4k+1$ and so $p=x^2+y^2$ with $x\equiv 1 \pmod 4$. In 1825, Gauss found the congruence

$$2x\equiv\binom{\frac{p-1}{2}}{\frac{p-1}{4}} \pmod p.$$

Similar congruences for $x \pmod p$ with $p=x^2+dy^2 (d\in\{2,3,5,7,11\})$ were found by Jacobi, Eisenstein and Cauchy, see [BEW]. For $d=3,7,11,19,43,67,163$ we know that for any prime p with $(\frac{p}{d})=1$, there are unique positive integers x and y such that $4p=x^2+dy^2$. In [R1,R2], [RPR], [PR], [PV] and [JM], the x was given by an appropriate cubic Jacobsthal sum. For example,

$$\sum_{x=0}^{p-1}\Big(\frac{x^3-96\cdot 11x+112\cdot 11^2}{p}\Big)=\begin{cases}(\frac{2}{p})(\frac{u}{11})u & \text{if } (\frac{p}{11})=1 \text{ and } 4p=u^2+11v^2,\\ 0 & \text{if } (\frac{p}{11})=-1.\end{cases}$$

In the paper, based on the work of Ishii[I], by considering the sums

$$\sum_{k=0}^{[p/6]}\binom{[\frac{p}{3}]}{k}\binom{[\frac{p}{6}]}{k}m^k,\ \sum_{k=0}^{[p/8]}\binom{[\frac{p}{8}]}{k}\binom{[\frac{3p}{8}]}{k}m^k,\ \sum_{k=0}^{p-1}\binom{-\frac{1}{3}}{k}^2m^k,\ \sum_{k=0}^{p-1}\binom{-\frac{1}{4}}{k}^2m^k$$

we establish many congruences for $x \pmod p$, where p is an odd prime and $x\in\mathbb{Z}$ is given by $p=x^2+dy^2$ or $4p=x^2+dy^2$. Here are four typical results:

$$\begin{aligned}
2x&\equiv\sum_{k=0}^{(p-1)/6}\binom{\frac{p-1}{3}}{k}\binom{\frac{p-1}{6}}{k}(-4)^k \pmod p\quad\text{for}\quad p=x^2+15y^2,\\
x&\equiv\sum_{k=0}^{(p-1)/6}\binom{\frac{p-1}{3}}{k}\binom{\frac{p-1}{6}}{k}\frac{1}{(-16)^k} \pmod p\quad\text{for}\quad 4p=x^2+51y^2,\\
2x&\equiv\sum_{k=0}^{(p-1)/8}\binom{\frac{p-1}{8}}{k}\binom{\frac{3(p-1)}{8}}{k}\frac{1}{(-882^2)^k} \pmod p\quad\text{for}\quad p=x^2+37y^2\equiv 1 \pmod 8,\\
2x&\equiv\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}{8}]}{k}\frac{1}{99^{4k}} \pmod p\quad\text{for}\quad p=x^2+58y^2.
\end{aligned}$$

We also pose many conjectures on congruences modulo p^2. For example, we conjecture that

$$y\equiv\frac{910}{9801}\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}{8}]}{k}\frac{1}{99^{4k}} \pmod p\quad\text{for}\quad p=2x^2+29y^2.$$

2. General congruences involving $\binom ak\binom{-\frac12-a}{k}$, $\binom ak\binom{a-\frac12}{k}$ and $\binom ak^2$.

For a prime p and $a\in\mathbb{Z}_p$ let $\langle a\rangle_p\in\{0,1,\ldots,p-1\}$ be given by $a\equiv\langle a\rangle_p \pmod p$. Let $P_n^{(\alpha,\beta)}(x)$ be the Jacobi polynomial defined by

$$P_n^{(\alpha,\beta)}(x)=\frac{1}{2^n}\sum_{k=0}^{n}\binom{n+\alpha}{k}\binom{n+\beta}{n-k}(x+1)^k(x-1)^{n-k}. \tag{2.1}$$

It is known that (see [AAR, p.315])

$$P_{2n}(x)=P_n^{(0,-\frac12)}(2x^2-1)\quad\text{and}\quad P_{2n+1}(x)=xP_n^{(0,\frac12)}(2x^2-1). \tag{2.2}$$

From [B, p.170] we know that

$$\begin{aligned}
P_n^{(\alpha,\beta)}(x)&=\binom{n+\alpha}{n}\sum_{k=0}^{\infty}\frac{(-n)_k(n+\alpha+\beta+1)_k}{(\alpha+1)_k\cdot k!}\Big(\frac{1-x}{2}\Big)^k\\
&=\binom{n+\alpha}{n}\sum_{k=0}^{n}\frac{\binom nk\binom{-n-\alpha-\beta-1}{k}}{\binom{-1-\alpha}{k}}\Big(\frac{x-1}{2}\Big)^k.
\end{aligned}$$

Thus,

$$P_n^{(0,\beta)}(x)=\sum_{k=0}^{n}\binom nk\binom{-n-\beta-1}{k}\Big(\frac{1-x}{2}\Big)^k. \tag{2.3}$$

Lemma 2.1. *Let n be a nonnegative integer. Then*

$$P_n(x)=x^n\sum_{k=0}^{[n/2]}\binom{n}{2k}\binom{-\frac12}{k}\Big(\frac{1}{x^2}-1\Big)^k,$$
$$P_{2n}(x)=\sum_{k=0}^{n}\binom{n}{k}\binom{-\frac12-n}{k}(1-x^2)^k,$$
$$P_{2n+1}(x)=x\sum_{k=0}^{n}\binom{n}{k}\binom{-\frac32-n}{k}(1-x^2)^k.$$

Proof. Since $\binom{-\frac12}{k}=\binom{2k}{k}\frac{1}{(-4)^k}$, by [G, (3.137)] we have

$$P_n(x)=x^n\sum_{k=0}^{[n/2]}\binom{n}{2k}\binom{2k}{k}\Big(\frac{x^2-1}{4x^2}\Big)^k=x^n\sum_{k=0}^{[n/2]}\binom{n}{2k}\binom{-\frac12}{k}\Big(\frac{1}{x^2}-1\Big)^k.$$

From (2.2) and (2.3) we see that

$$\begin{aligned}P_{2n}(x)=P_n^{(0,-\frac12)}(2x^2-1)&=\sum_{k=0}^{n}\binom{n}{k}\binom{-n+\frac12-1}{k}\Big(\frac{1-(2x^2-1)}{2}\Big)^k\\&=\sum_{k=0}^{n}\binom{n}{k}\binom{-\frac12-n}{k}(1-x^2)^k\end{aligned}$$

and

$$\begin{aligned}P_{2n+1}(x)=xP_n^{(0,\frac12)}(2x^2-1)&=x\sum_{k=0}^{n}\binom{n}{k}\binom{-n-\frac12-1}{k}\Big(\frac{1-(2x^2-1)}{2}\Big)^k\\&=x\sum_{k=0}^{n}\binom{n}{k}\binom{-\frac32-n}{k}(1-x^2)^k.\end{aligned}$$

This proves the lemma.

Theorem 2.1. *Let p be an odd prime and let m be a positive integer such that $p\nmid m$. Then*

$$\begin{aligned}&\sum_{k=0}^{[\frac{p}{2m}]}\binom{[\frac{p}{2m}]}{k}\binom{[\frac{(m-1)p}{2m}]}{k}(1-t)^k\\&\equiv t^{[\frac{p}{2m}]}\sum_{k=0}^{[\frac{p}{2m}]}\binom{[\frac{p}{m}]}{2k}\binom{\frac{p-1}{2}}{k}\Big(\frac1t-1\Big)^k\\&\equiv\begin{cases}P_{[\frac{p}{m}]}(\sqrt t)\pmod p & \text{if } 2\mid [\frac{p}{m}],\\ P_{[\frac{p}{m}]}(\sqrt t)/\sqrt t\pmod p & \text{if } 2\nmid [\frac{p}{m}].\end{cases}\end{aligned}$$

Proof. Suppose $p = 2mk + r$ with $k \in \mathbb{Z}$ and $r \in \{0, 1, \dots, 2m-1\}$. Then $[\frac{p}{m}] = [\frac{r}{m}]$ and so $2 \mid [\frac{p}{m}]$ if and only if $r < m$. Hence

$$\begin{aligned}
&\Big[\frac{p}{2m}\Big] + \Big[\frac{(m-1)p}{2m}\Big] \\
&= k + k(m-1) + \Big[\frac{(m-1)r}{2m}\Big] = \frac{p-r}{2} + \Big[\frac{(m-1)r}{2m}\Big] \\
&= \frac{p-1}{2} + \Big[\frac{(m-1)r + m(1-r)}{2m}\Big] = \frac{p-1}{2} + \Big[\frac{m-r}{2m}\Big] \\
&= \begin{cases} \frac{p-1}{2} & \text{if } 2 \mid [\frac{p}{m}], \\ \frac{p-3}{2} & \text{if } 2 \nmid [\frac{p}{m}]. \end{cases}
\end{aligned}$$

Thus, if $2 \mid [\frac{p}{m}]$, using Lemma 2.1 and the above we get

$$\begin{aligned}
P_{[\frac{p}{m}]}(\sqrt{t}) = P_{2[\frac{p}{2m}]}(\sqrt{t}) &= \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{2m}]}{k} \binom{-\frac{1}{2} - [\frac{p}{2m}]}{k} (1-t)^k \\
&\equiv \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{2m}]}{k} \binom{\frac{p-1}{2} - [\frac{p}{2m}]}{k} (1-t)^k \\
&= \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{2m}]}{k} \binom{[\frac{(m-1)p}{2m}]}{k} (1-t)^k \pmod p.
\end{aligned}$$

If $2 \nmid [\frac{p}{m}]$, using Lemma 2.1 and the above we get

$$\begin{aligned}
P_{[\frac{p}{m}]}(\sqrt{t})/\sqrt{t} = P_{2[\frac{p}{2m}]+1}(\sqrt{t})/\sqrt{t} &= \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{2m}]}{k} \binom{-\frac{3}{2} - [\frac{p}{2m}]}{k} (1-t)^k \\
&\equiv \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{2m}]}{k} \binom{\frac{p-3}{2} - [\frac{p}{2m}]}{k} (1-t)^k \\
&= \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{2m}]}{k} \binom{[\frac{(m-1)p}{2m}]}{k} (1-t)^k \pmod p.
\end{aligned}$$

By Lemma 2.1, we also have

$$\begin{aligned}
&t^{[\frac{p}{2m}]} \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{m}]}{2k} \binom{\frac{p-1}{2}}{k} \Big(\frac{1}{t} - 1\Big)^k \\
&\equiv (\sqrt{t})^{2[\frac{p}{2m}]} \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{m}]}{2k} \binom{-\frac{1}{2}}{k} \Big(\frac{1}{t} - 1\Big)^k \\
&= \begin{cases} (\sqrt{t})^{[\frac{p}{m}]} \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{m}]}{2k} \binom{-\frac{1}{2}}{k} (\frac{1}{t} - 1)^k = P_{[\frac{p}{m}]}(\sqrt{t}) \pmod p & \text{if } 2 \mid [\frac{p}{m}], \\ (\sqrt{t})^{[\frac{p}{m}]-1} \sum_{k=0}^{[\frac{p}{2m}]} \binom{[\frac{p}{m}]}{2k} \binom{-\frac{1}{2}}{k} (\frac{1}{t} - 1)^k = \frac{P_{[\frac{p}{m}]}(\sqrt{t})}{\sqrt{t}} \pmod p & \text{if } 2 \nmid [\frac{p}{m}]. \end{cases}
\end{aligned}$$

This completes the proof.

Lemma 2.2. *Let p be an odd prime and $m \in \{1, 2, \dots, \frac{p-1}{2}\}$. Then $P_{p-1-m}(x) \equiv P_m(x)$.*

Proof. Since $m < \frac{p}{2}$ we have $p - 1 - m \geq m$. Note that $\binom{-t}{k} = (-1)^k \binom{t+k-1}{k}$. By using (1.3) we have

$$\begin{aligned}
P_{p-1-m}(x) &= \sum_{k=0}^{p-1-m} \binom{p-1-m}{k} \binom{p-1-m+k}{k} \Big(\frac{x-1}{2}\Big)^k \\
&\equiv \sum_{k=0}^{p-1-m} \binom{-1-m}{k} \binom{-1-m+k}{k} \Big(\frac{x-1}{2}\Big)^k \\
&= \sum_{k=0}^{p-1-m} (-1)^k \binom{m+k}{k} \cdot (-1)^k \binom{m}{k} \Big(\frac{x-1}{2}\Big)^k \\
&= \sum_{k=0}^{m} \binom{m}{k} \binom{m+k}{k} \Big(\frac{x-1}{2}\Big)^k = P_m(x) \pmod p.
\end{aligned}$$

This proves the lemma.

Lemma 2.3. *Let p be an odd prime and $m \in \{0, 1, \dots, p-1\}$. Then*

$$P_{p+m}(x) \equiv x^p P_m(x) \pmod p.$$

Proof. If $a_i, b_i \in \{0, 1, \dots, p-1\}$, then Lucas theorem asserts that

$$\binom{a_0 + a_1 p + \cdots + a_n p^n}{b_0 + b_1 p + \cdots + b_n p^n} \equiv \binom{a_0}{b_0} \binom{a_1}{b_1} \cdots \binom{a_n}{b_n} \pmod p.$$

Thus, for $0 \leq r \leq m < p$ we have

$$\binom{2p+m+r}{p+r} \equiv \begin{cases} \binom{2}{1}\binom{m+r}{r} \pmod p & \text{if } r < p-m, \\ \binom{3p}{p+r} \equiv \binom{3}{1}\binom{0}{r} = 0 \equiv \binom{2}{1}\binom{m+r}{r} \pmod p & \text{if } r = p-m. \\ \binom{3}{1}\binom{m+r-p}{r} = 0 \equiv \binom{2}{1}\binom{m+r}{r} \pmod p & \text{if } r > p-m. \end{cases}$$

Hence, using (1.3) we see that

$$\begin{aligned}
P_{p+m}(x) &= \sum_{k=0}^{p+m}\binom{p+m}{k}\binom{p+m+k}{k}\Big(\frac{x-1}{2}\Big)^k \\
&= \sum_{k=0}^{p-1}\binom{p+m}{k}\binom{p+m+k}{k}\Big(\frac{x-1}{2}\Big)^k \\
&\quad + \sum_{k=p}^{p+m}\binom{p+m}{k}\binom{p+m+k}{k}\Big(\frac{x-1}{2}\Big)^k \\
&\equiv \sum_{k=0}^{p-1}\binom{m}{k}\binom{m+k}{k}\Big(\frac{x-1}{2}\Big)^k \\
&\quad + \sum_{r=0}^{m}\binom{p+m}{p+r}\binom{2p+m+r}{p+r}\Big(\frac{x-1}{2}\Big)^{p+r} \\
&\equiv \sum_{k=0}^{m}\binom{m}{k}\binom{m+k}{k}\Big(\frac{x-1}{2}\Big)^k + \sum_{r=0}^{m}\binom{m}{r}\cdot 2\binom{m+r}{r}\Big(\frac{x-1}{2}\Big)^{p+r} \\
&\equiv (1+(x-1)^p)\sum_{k=0}^{m}\binom{m}{k}\binom{m+k}{k}\Big(\frac{x-1}{2}\Big)^k \\
&\equiv x^p P_m(x) \pmod p.
\end{aligned}$$

Thus the lemma is proved.

Theorem 2.2. *Let p be an odd prime and $a, t \in \mathbb{Z}_p$ with $t \not\equiv 0 \pmod p$. Then*

$$\begin{aligned}
P_{2\langle a\rangle_p}(\sqrt{t}) &\equiv \sum_{k=0}^{p-1}\binom{a}{k}\binom{-\frac{1}{2}-a}{k}(1-t)^k \\
&\equiv t^{\langle a\rangle_p}\sum_{k=0}^{p-1}\binom{a}{k}\binom{a-\frac{1}{2}}{k}\Big(1-\frac{1}{t}\Big)^k \\
&\equiv \Big(\frac{t}{p}\Big)t^{[\frac{2\langle a\rangle_p}{p}]}\sum_{k=0}^{p-1}\binom{-1-a}{k}\binom{a-\frac{1}{2}}{k}(1-t)^k \\
&\equiv \Big(\frac{t}{p}\Big)t^{[\frac{2\langle a\rangle_p}{p}]-\langle a\rangle_p}\sum_{k=0}^{p-1}\binom{-1-a}{k}\binom{-\frac{1}{2}-a}{k}\Big(1-\frac{1}{t}\Big)^k \pmod p.
\end{aligned}$$

Proof. For $\beta \in \mathbb{Z}_p$ we have

$$\begin{aligned}
&\sum_{k=0}^{p-1}\binom{a}{k}\binom{a+\beta}{k}\Big(\frac{t-1}{t}\Big)^k \\
&\equiv \sum_{k=0}^{p-1}\binom{\langle a\rangle_p}{k}\binom{\langle a\rangle_p+\beta}{k}\Big(\frac{t-1}{t}\Big)^k \\
&= \sum_{k=0}^{\langle a\rangle_p}\binom{\langle a\rangle_p}{k}\binom{\langle a\rangle_p+\beta}{k}\Big(\frac{t-1}{t}\Big)^k \\
&= \sum_{k=0}^{\langle a\rangle_p}\binom{\langle a\rangle_p}{\langle a\rangle_p-k}\binom{\langle a\rangle_p+\beta}{\langle a\rangle_p-k}\Big(\frac{t-1}{t}\Big)^{\langle a\rangle_p-k} \\
&= \frac{1}{(2t)^{\langle a\rangle_p}}\sum_{k=0}^{\langle a\rangle_p}\binom{\langle a\rangle_p}{k}\binom{\langle a\rangle_p+\beta}{\langle a\rangle_p-k}((2t-1)+1)^k((2t-1)-1)^{\langle a\rangle_p-k} \\
&= \frac{1}{t^{\langle a\rangle_p}}P_{\langle a\rangle_p}^{(0,\beta)}(2t-1) \pmod p.
\end{aligned}$$

Thus, by (2.2) and the fact $\langle -1-a\rangle_p = p-1-\langle a\rangle_p$ we get

$$\sum_{k=0}^{p-1}\binom{a}{k}\binom{a-\frac12}{k}\Big(\frac{t-1}{t}\Big)^k \equiv \frac{1}{t^{\langle a\rangle_p}}P_{\langle a\rangle_p}^{(0,-\frac12)}(2t-1) = \frac{1}{t^{\langle a\rangle_p}}P_{2\langle a\rangle_p}(\sqrt{t}) \pmod p$$

and

$$\begin{aligned}
&\sum_{k=0}^{p-1}\binom{-1-a}{k}\binom{-\frac12-a}{k}\Big(\frac{t-1}{t}\Big)^k \\
&\equiv t^{-\langle -1-a\rangle_p}P_{\langle -1-a\rangle_p}^{(0,\frac12)}(2t-1) = t^{-\langle -1-a\rangle_p}P_{2\langle -1-a\rangle_p+1}(\sqrt{t})/\sqrt{t} \\
&\equiv t^{\langle a\rangle_p}P_{2(p-1-\langle a\rangle_p)+1}(\sqrt{t})/\sqrt{t} \pmod p.
\end{aligned}$$

Using (2.3) and (2.2) we see that

$$\begin{aligned}
&\sum_{k=0}^{p-1}\binom{a}{k}\binom{-\frac12-a}{k}(1-t)^k \\
&\equiv \sum_{k=0}^{p-1}\binom{\langle a\rangle_p}{k}\binom{-\frac12-\langle a\rangle_p}{k}(1-t)^k = \sum_{k=0}^{\langle a\rangle_p}\binom{\langle a\rangle_p}{k}\binom{-\frac12-\langle a\rangle_p}{k}(1-t)^k \\
&= P_{\langle a\rangle_p}^{(0,-\frac12)}(2t-1) = P_{2\langle a\rangle_p}(\sqrt{t}) \pmod p
\end{aligned}$$

and

$$\begin{aligned}
&\sum_{k=0}^{p-1}\binom{-1-a}{k}\binom{a-\frac{1}{2}}{k}(1-t)^k\\
&\equiv\sum_{k=0}^{p-1}\binom{p-1-\langle a\rangle_p}{k}\binom{1+a-\frac{1}{2}-1}{k}(1-t)^k\\
&\equiv\sum_{k=0}^{p-1-\langle a\rangle_p}\binom{p-1-\langle a\rangle_p}{k}\binom{-(p-1-\langle a\rangle_p)-\frac{1}{2}-1}{k}(1-t)^k\\
&=P_{p-1-\langle a\rangle_p}^{(0,\frac{1}{2})}(2t-1)=P_{2(p-1-\langle a\rangle_p)+1}(\sqrt{t})/\sqrt{t}\ (\mathrm{mod}\ p).
\end{aligned}$$

To complete the proof, using Lemmas 2.2 and 2.3 we note that

$$\begin{aligned}
&P_{2(p-1-\langle a\rangle_p)+1}(\sqrt{t})/\sqrt{t}\\
&\equiv\begin{cases}\frac{P_{p-1-2\langle a\rangle_p}(\sqrt{t})(\sqrt{t})^p}{\sqrt{t}}\equiv(\frac{t}{p})P_{2\langle a\rangle_p}(\sqrt{t})\ (\mathrm{mod}\ p) & \text{if } \langle a\rangle_p<\frac{p}{2},\\ \frac{P_{p-1-(2\langle a\rangle_p-p)}(\sqrt{t})}{\sqrt{t}}\equiv\frac{P_{2\langle a\rangle_p-p}(\sqrt{t})}{\sqrt{t}}\equiv\frac{P_{2\langle a\rangle_p}(\sqrt{t})}{(\sqrt{t})^{p+1}} & \\ \quad\equiv(\frac{t}{p})\frac{1}{t}P_{2\langle a\rangle_p}(\sqrt{t})\ (\mathrm{mod}\ p) & \text{if } \langle a\rangle_p>\frac{p}{2}.\end{cases}
\end{aligned}$$

Theorem 2.3. *Let p be an odd prime and $a,t\in\mathbb{Z}_p$ with $t\not\equiv 0,1\ (\mathrm{mod}\ p)$. Then*

$$\begin{aligned}
\sum_{k=0}^{p-1}\binom{a}{k}^2t^k&\equiv t^{\langle a\rangle_p}\sum_{k=0}^{p-1}\frac{\binom{a}{k}^2}{t^k}\equiv(t-1)^{\langle a\rangle_p}P_{\langle a\rangle_p}\Big(\frac{t+1}{t-1}\Big)\\
&\equiv(t-1)^{\langle a\rangle_p}\sum_{k=0}^{p-1}\binom{a}{k}\binom{-1-a}{k}\frac{1}{(1-t)^k}\\
&\equiv(t-1)^{2\langle a\rangle_p}\sum_{k=0}^{p-1}\binom{-1-a}{k}^2t^k\ (\mathrm{mod}\ p).
\end{aligned}$$

Proof. It is clear that

$$\begin{aligned}
\sum_{k=0}^{p-1}\binom{a}{k}^2t^k&\equiv\sum_{k=0}^{\langle a\rangle_p}\binom{\langle a\rangle_p}{k}^2t^k=\sum_{s=0}^{\langle a\rangle_p}\binom{\langle a\rangle_p}{\langle a\rangle_p-s}^2t^{\langle a\rangle_p-s}\\
&=t^{\langle a\rangle_p}\sum_{k=0}^{\langle a\rangle_p}\binom{\langle a\rangle_p}{k}^2t^{-k}\equiv t^{\langle a\rangle_p}\sum_{k=0}^{p-1}\frac{\binom{a}{k}^2}{t^k}\ (\mathrm{mod}\ p).
\end{aligned}$$

Using (2.1) we see that for $x \neq 1$,

$$\begin{aligned} P_{\langle a\rangle_p}(x) = P^{(0,0)}_{\langle a\rangle_p}(x) &= \Big(\frac{x-1}{2}\Big)^{\langle a\rangle_p} \sum_{k=0}^{\langle a\rangle_p} \binom{\langle a\rangle_p}{k}^2 \Big(\frac{x+1}{x-1}\Big)^k \\ &= \Big(\frac{x-1}{2}\Big)^{\langle a\rangle_p} \sum_{k=0}^{p-1} \binom{\langle a\rangle_p}{k}^2 \Big(\frac{x+1}{x-1}\Big)^k \\ &\equiv \Big(\frac{x-1}{2}\Big)^{\langle a\rangle_p} \sum_{k=0}^{p-1} \binom{a}{k}^2 \Big(\frac{x+1}{x-1}\Big)^k \pmod p. \end{aligned}$$

Set $x = \frac{t+1}{t-1}$. Then $t = \frac{x+1}{x-1}$ and $\frac{x-1}{2} = \frac{1}{t-1}$. Now substituting x with $\frac{t+1}{t-1}$ in the above congruence we obtain

$$\sum_{k=0}^{p-1} \binom{a}{k}^2 t^k \equiv (t-1)^{\langle a\rangle_p} P_{\langle a\rangle_p}\Big(\frac{t+1}{t-1}\Big) \pmod p.$$

Clearly $\langle -1-a\rangle_p = p-1-\langle a\rangle_p$. Thus, using Lemma 2.2 and the above we see that

$$\begin{aligned} &\sum_{k=0}^{p-1} \binom{-1-a}{k}^2 t^k \\ &\equiv (t-1)^{\langle -1-a\rangle_p} P_{\langle -1-a\rangle_p}\Big(\frac{t+1}{t-1}\Big) = (t-1)^{p-1-\langle a\rangle_p} P_{p-1-\langle a\rangle_p}\Big(\frac{t+1}{t-1}\Big) \\ &\equiv (t-1)^{-\langle a\rangle_p} P_{\langle a\rangle_p}\Big(\frac{t+1}{t-1}\Big) \equiv (t-1)^{-2\langle a\rangle_p} \sum_{k=0}^{p-1} \binom{a}{k}^2 t^k \pmod p. \end{aligned}$$

To complete the proof, using (2.3) we note that

$$\begin{aligned} P_{\langle a\rangle_p}\Big(\frac{t+1}{t-1}\Big) &= \sum_{k=0}^{\langle a\rangle_p} \binom{\langle a\rangle_p}{k}\binom{-1-\langle a\rangle_p}{k} \frac{1}{(1-t)^k} \\ &= \sum_{k=0}^{p-1} \binom{\langle a\rangle_p}{k}\binom{-1-\langle a\rangle_p}{k} \frac{1}{(1-t)^k} \\ &\equiv \sum_{k=0}^{p-1} \binom{a}{k}\binom{-1-a}{k} \frac{1}{(1-t)^k} \pmod p. \end{aligned}$$

3. Congruences for $\sum_{x=0}^{p-1}(\frac{x^3+mx+n}{p}) \pmod p$.

Theorem 3.1. *Let p be an odd prime and let $t \in \mathbb{Z}_p$. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{5}{12}}{k}(1-t)^k\\&\equiv t^{\langle -\frac{1}{12}\rangle_p}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{7}{12}}{k}\Big(1-\frac{1}{t}\Big)^k\\&\equiv\begin{cases}P_{[\frac{p}{6}]}(\sqrt{t}) \pmod{p} & \text{if } p\equiv 1 \pmod 4,\\ (\frac{t}{p})P_{[\frac{p}{6}]}(\sqrt{t})\sqrt{t} \pmod{p} & \text{if } p\equiv 3 \pmod 4.\end{cases}\end{aligned}$$

Proof. Taking $a=-\frac{1}{12}$ in Theorem 2.2 and then applying Lemmas 2.2-2.3 we see that

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{5}{12}}{k}(1-t)^k\\&\equiv t^{\langle -\frac{1}{12}\rangle_p}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{7}{12}}{k}\Big(\frac{t-1}{t}\Big)^k\equiv P_{2\langle -\frac{1}{12}\rangle_p}(\sqrt{t})\\&=\begin{cases}P_{\frac{p-1}{6}}(\sqrt{t}) \pmod{p} & \text{if } 12\mid p-1,\\ P_{\frac{5p-1}{6}}(\sqrt{t})\equiv P_{\frac{p-5}{6}}(\sqrt{t}) \pmod{p} & \text{if } 12\mid p-5,\\ P_{\frac{7p-1}{6}}(\sqrt{t})\equiv(\sqrt{t})^pP_{\frac{p-1}{6}}(\sqrt{t}) \pmod{p} & \text{if } 12\mid p-7,\\ P_{\frac{11p-1}{6}}(\sqrt{t})\equiv(\sqrt{t})^pP_{\frac{5p-1}{6}}(\sqrt{t})\equiv(\sqrt{t})^pP_{\frac{p-5}{6}}(\sqrt{t}) \pmod{p} & \text{if } 12\mid p-11.\end{cases}\end{aligned}$$

To see the result, we note that $(\sqrt{t})^p=t^{\frac{p-1}{2}}\sqrt{t}\equiv(\frac{t}{p})\sqrt{t} \pmod{p}$.

Lemma 3.1 ([S5, Theorem 2.2]). *Let $p>3$ be a prime and $m,n\in\mathbb{Z}_p$ with $m\not\equiv 0 \pmod{p}$. Then*

$$\sum_{x=0}^{p-1}\Big(\frac{x^3+mx+n}{p}\Big)\equiv\begin{cases}-(-3m)^{\frac{p-1}{4}}P_{[\frac{p}{6}]}\big(\frac{3n\sqrt{-3m}}{2m^2}\big) \pmod{p} & \text{if } p\equiv 1 \pmod 4,\\ -\frac{(-3m)^{\frac{p+1}{4}}}{\sqrt{-3m}}P_{[\frac{p}{6}]}\big(\frac{3n\sqrt{-3m}}{2m^2}\big) \pmod{p} & \text{if } p\equiv 3 \pmod 4.\end{cases}$$

Theorem 3.2. *Let $p>3$ be a prime and $m,n\in\mathbb{Z}_p$ with $mn\not\equiv 0 \pmod{p}$. Then*

$$\begin{aligned}&\sum_{x=0}^{p-1}\Big(\frac{x^3+mx+n}{p}\Big)\\&\equiv\begin{cases}-(-3m)^{\frac{p-1}{4}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{5}{12}}{k}\big(\frac{4m^3+27n^2}{4m^3}\big)^k \pmod{p} & \text{if } 4\mid p-1,\\ \frac{2m}{9n}\big(\frac{-3m}{p}\big)(-3m)^{\frac{p+1}{4}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{5}{12}}{k}\big(\frac{4m^3+27n^2}{4m^3}\big)^k \pmod{p} & \text{if } 4\mid p-3.\end{cases}\\&\equiv\begin{cases}(-1)^{\frac{p+1}{2}}\big(\frac{n}{2}\big)^{\frac{p-1}{6}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{7}{12}}{k}\big(\frac{4m^3+27n^2}{27n^2}\big)^k \pmod{p} & \text{if } 3\mid p-1,\\ (-1)^{\frac{p+1}{2}}\frac{3}{m}\big(\frac{2}{n}\big)^{\frac{p-5}{6}}\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{7}{12}}{k}\big(\frac{4m^3+27n^2}{27n^2}\big)^k \pmod{p} & \text{if } 3\mid p-2.\end{cases}\end{aligned}$$

Proof. Set $t=-\frac{27n^2}{4m^3}$. Then $1-t=\frac{4m^3+27n^2}{4m^3}$. Since

$$t^{\langle -\frac{1}{12}\rangle_p}=\begin{cases}(-\frac{27n^2}{4m^3})^{\frac{p-1}{12}}=(-\frac{3}{m})^{\frac{p-1}{4}}(\frac{n}{2})^{\frac{p-1}{6}} & \text{if } p\equiv 1 \pmod{12},\\ (-\frac{27n^2}{4m^3})^{\frac{5p-1}{12}}\equiv(\frac{-3m}{p})(-\frac{m}{3})^{\frac{p-5}{4}}(\frac{2}{n})^{\frac{p-5}{6}} \pmod p & \text{if } p\equiv 5 \pmod{12},\\ (-\frac{27n^2}{4m^3})^{\frac{7p-1}{12}}\equiv(\frac{-3m}{p})(-\frac{3}{m})^{\frac{p+5}{4}}(\frac{n}{2})^{\frac{p+5}{6}} \pmod p & \text{if } p\equiv 7 \pmod{12},\\ (-\frac{27n^2}{4m^3})^{\frac{11p-1}{12}}\equiv(-\frac{m}{3})^{\frac{p-11}{4}}(\frac{2}{n})^{\frac{p-11}{6}} \pmod p & \text{if } p\equiv 11 \pmod{12},\end{cases}$$

using Theorem 3.1 and Lemma 3.1 we deduce the result.

Corollary 3.1. *Let p be a prime with $p\neq 2,3,19$. Then*

$$\sum_{k=0}^{p-1}\binom{-\frac{1}{12}}{k}\binom{-\frac{7}{12}}{k}\frac{1}{513^k}$$
$$\equiv\begin{cases}0 \pmod p & \textit{if } (\frac{p}{19})=-1,\\ -19^{-\frac{p-1}{3}}(\frac{-2}{p})(\frac{u}{19})u \pmod p & \textit{if } (\frac{p}{19})=(\frac{p}{3})=1 \textit{ and so } 4p=u^2+19v^2.\\ \frac{8}{3}\cdot 19^{\frac{p-2}{3}}(\frac{-2}{p})(\frac{u}{19})u \pmod p & \textit{if } (\frac{p}{19})=-(\frac{p}{3})=1 \textit{ and so } 4p=u^2+19v^2.\end{cases}$$

Proof. From [RPR], [JM] and [PV] we know that

$$(3.1)\quad \sum_{x=0}^{p-1}\Big(\frac{x^3-8\cdot 19x+2\cdot 19^2}{p}\Big)=\begin{cases}(\frac{2}{p})(\frac{u}{19})u & \text{if } (\frac{p}{19})=1 \text{ and } 4p=u^2+19v^2,\\ 0 & \text{if } (\frac{p}{19})=-1.\end{cases}$$

Thus, taking $m=-8\cdot 19$ and $n=2\cdot 19^2$ in Theorem 3.2 we deduce the result.

For a simpler congruence for $u \pmod p$ with $4p=u^2+19v^2$, see [LH, p.269].

Theorem 3.3. *Let $p>3$ be a prime and $m,n\in\mathbb{Z}_p$ with $mn\not\equiv 0 \pmod p$. Then*

$$\sum_{x=0}^{p-1}\Big(\frac{x^3+mx+n}{p}\Big)$$
$$\equiv\begin{cases}-(-3m)^{\frac{p-1}{4}}\sum_{k=0}^{[p/12]}\binom{[\frac{p}{12}]}{k}\binom{[\frac{5p}{12}]}{k}(\frac{4m^3+27n^2}{4m^3})^k \pmod p & \textit{if } 4\mid p-1,\\ -\frac{3n}{2m^2}(-3m)^{\frac{p+1}{4}}\sum_{k=0}^{[p/12]}\binom{[\frac{p}{12}]}{k}\binom{[\frac{5p}{12}]}{k}(\frac{4m^3+27n^2}{4m^3})^k \pmod p & \textit{if } 4\mid p-3\end{cases}$$
$$\equiv\begin{cases}(-1)^{\frac{p+1}{2}}(\frac{n}{2})^{\frac{p-1}{6}}\sum_{k=0}^{[p/12]}\binom{[\frac{p}{6}]}{2k}\binom{\frac{p-1}{2}}{k}(-\frac{4m^3+27n^2}{27n^2})^k \pmod p & \textit{if } 3\mid p-1,\\ (-1)^{\frac{p+1}{2}}\frac{m}{3}(\frac{n}{2})^{\frac{p-5}{6}}\sum_{k=0}^{[p/12]}\binom{[\frac{p}{6}]}{2k}\binom{\frac{p-1}{2}}{k}(-\frac{4m^3+27n^2}{27n^2})^k \pmod p & \textit{if } 3\mid p-2.\end{cases}$$

Proof. Putting $m=6$ and $t=-\frac{27n^2}{4m^3}$ in Theorem 2.1 we see that

$$\begin{aligned}&\sum_{k=0}^{[p/12]}\binom{[\frac{p}{12}]}{k}\binom{[\frac{5p}{12}]}{k}\Big(\frac{4m^3+27n^2}{4m^3}\Big)^k\\&\equiv\Big(-\frac{27n^2}{4m^3}\Big)^{[\frac{p}{12}]}\sum_{k=0}^{[p/12]}\binom{[\frac{p}{6}]}{2k}\binom{\frac{p-1}{2}}{k}\Big(-\frac{4m^3+27n^2}{27n^2}\Big)^k\\&\equiv\begin{cases}P_{[\frac{p}{6}]}(\frac{3n\sqrt{-3m}}{2m^2})\pmod p & \text{if } p\equiv 1,5\pmod{12},\\ P_{[\frac{p}{6}]}(\frac{3n\sqrt{-3m}}{2m^2})/(\frac{3n\sqrt{-3m}}{2m^2})\pmod p & \text{if } p\equiv 7,11\pmod{12}.\end{cases}\end{aligned}$$

Now applying Lemma 3.1 we deduce the result.

Let $p>3$ be a prime and $S_p(m,n)=\sum_{x=0}^{p-1}(\frac{x^3+mx+n}{p})$. It is known that (see for example [S5, S6], [R1,R2], [RPR], [JM], [PV] and [W])

$$S_p(-11,14)=\begin{cases}(-1)^{\frac{p+3}{4}}2a & \text{if } 4\mid p-1,\ p=a^2+b^2 \text{ and } 4\mid a-1,\\ 0 & \text{if } p\equiv 3\pmod 4,\end{cases}\tag{3.2}$$

$$S_p(-30,56)=\begin{cases}(-1)^{\frac{p+7}{8}}(\frac{3}{p})2c & \text{if } p\equiv 1\pmod 8,\ p=c^2+2d^2 \text{ and } 4\mid c-1,\\ (-1)^{\frac{p-3}{8}}(\frac{3}{p})2c & \text{if } p\equiv 3\pmod 8,\ p=c^2+2d^2 \text{ and } 4\mid c-1,\\ 0 & \text{if } p\equiv 5,7\pmod 8,\end{cases}\tag{3.3}$$

$$S_p(-15,22)=\begin{cases}-2A & \text{if } 3\mid p-1,\ p=A^2+3B^2 \text{ and } 3\mid A-1,\\ 0 & \text{if } p\equiv 2\pmod 3,\end{cases}\tag{3.4}$$

$$S_p(-120,506)=\begin{cases}(\frac{2}{p})L & \text{if } 3\mid p-1,\ 4p=L^2+27M^2 \text{ and } 3\mid L-1,\\ 0 & \text{if } p\equiv 2\pmod 3,\end{cases}\tag{3.5}$$

$$S_p(-35,98)=\begin{cases}(-1)^{\frac{p+1}{2}}2(\frac{C}{7})C & \text{if } p\equiv 1,2,4\pmod 7 \text{ and } p=C^2+7D^2,\\ 0 & \text{if } p\equiv 3,5,6\pmod 7,\end{cases}\tag{3.6}$$

$$S_p(-595,5586)=\begin{cases}(-1)^{\frac{p+1}{2}}2C(\frac{C}{7}) & \text{if } p=C^2+7D^2\equiv 1,2,4\pmod 7,\\ 0 & \text{if } p\equiv 3,5,6\pmod 7,\end{cases}\tag{3.7}$$

$$S_p(-96\cdot 11,112\cdot 11^2)=\begin{cases}(\frac{2}{p})(\frac{u}{11})u & \text{if } (\frac{p}{11})=1 \text{ and } 4p=u^2+11v^2,\\ 0 & \text{if } (\frac{p}{11})=-1.\end{cases}\tag{3.8}$$

$$S_p(-80\cdot 43, 42\cdot 43^2)=\begin{cases}(\frac 2p)(\frac u{43})u & \text{if } (\frac p{43})=1 \text{ and } 4p=u^2+43v^2,\\ 0 & \text{if } (\frac p{43})=-1,\end{cases}\tag{3.9}$$

$$S_p(-440\cdot 67, 434\cdot 67^2)=\begin{cases}(\frac 2p)(\frac u{67})u & \text{if } (\frac p{67})=1 \text{ and } 4p=u^2+67v^2,\\ 0 & \text{if } (\frac p{67})=-1,\end{cases}\tag{3.10}$$

$$\begin{aligned}&S_p(-80\cdot 23\cdot 29\cdot 163, 14\cdot 11\cdot 19\cdot 127\cdot 163^2)\\ &=\begin{cases}(\frac 2p)(\frac u{163})u & \text{if } (\frac p{163})=1 \text{ and } 4p=u^2+163v^2,\\ 0 & \text{if } (\frac p{163})=-1.\end{cases}\end{aligned}\tag{3.11}$$

Using (3.2)-(3.11) and Theorems 3.2 and 3.3 we may deduce the results similar to Corollary 3.1.

4. Congruences for $\sum_{k=0}^{[p/6]}\binom{[\frac p3]}k\binom{[\frac p6]}k(1-t)^k \pmod p$.

Theorem 4.1. *Let p be a prime with $p\equiv 1,4 \pmod 5$. Then*

$$\begin{aligned}&\sum_{k=0}^{[p/6]}\binom{[\frac p3]}k\binom{[\frac p6]}k(-4)^k\\ &\equiv\begin{cases}2x \pmod p & \text{if } p\equiv 1,4 \pmod{15} \text{ and so } p=x^2+15y^2 \text{ with } 3\mid x-1,\\ 0 \pmod p & \text{if } p\equiv 11,14 \pmod{15}.\end{cases}\end{aligned}$$

Proof. Taking $m=3$ and $t=5$ in Theorem 2.1 we get

$$\sum_{k=0}^{[p/6]}\binom{[\frac p3]}k\binom{[\frac p6]}k(-4)^k\equiv\begin{cases}P_{[\frac p3]}(\sqrt5) \pmod p & \text{if } p\equiv 1 \pmod 3,\\ P_{\frac{p-2}3}(\sqrt5)/\sqrt5 \pmod p & \text{if } p\equiv 2 \pmod 3.\end{cases}$$

From [S6, Theorem 4.6] we know that

$$P_{[\frac p3]}(\sqrt5)\equiv\begin{cases}2x \pmod p & \text{if } p=x^2+15y^2\equiv 1,4 \pmod{15} \text{ and } 3\mid x-1,\\ 0 \pmod p & \text{if } p\equiv 11,14 \pmod{15}.\end{cases}$$

Thus the result follows.

Conjecture 4.1. *Let $p>5$ be a prime. Then*

$$\begin{aligned}\sum_{k=0}^{p-1}\binom{-\frac13}k\binom{-\frac16}k(-4)^k&\equiv\Big(\frac 5p\Big)5^{\frac{1-(\frac p3)}2}\sum_{k=0}^{p-1}\binom{-\frac23}k\binom{-\frac56}k(-4)^k\\ &\equiv\begin{cases}(\frac x3)(2x-\frac p{2x}) \pmod{p^2} & \text{if } p=x^2+15y^2,\\ -(\frac x3)(10x-\frac p{2x}) \pmod{p^2} & \text{if } p=5x^2+3y^2,\\ 0 \pmod p & \text{if } p\equiv 17,23 \pmod{30}\end{cases}\end{aligned}$$

and so

$$2x\Big(\frac x3\Big)\equiv\sum_{k=0}^{(p-5)/6}\binom{\frac{p-2}3}k\binom{\frac{p-5}6}k(-4)^k \pmod p \quad \text{for} \quad p=5x^2+3y^2.$$

Theorem 4.2. *Let p be a prime such that $p\equiv 1,7 \pmod 8$. Then*

$$2^{[\frac{p}{3}]}\sum_{k=0}^{[p/6]}\binom{[\frac{p}{3}]}{2k}\binom{\frac{p-1}{2}}{k}\equiv\sum_{k=0}^{[p/6]}\binom{[\frac{p}{3}]}{k}\binom{[\frac{p}{6}]}{k}\frac{1}{2^k}$$
$$\equiv\begin{cases}2x \pmod p & \text{if } p\equiv 1,7 \pmod{24} \text{ and so } p=x^2+6y^2 \text{ with } 3\mid x-1,\\ 0 \pmod p & \text{if } p\equiv 17,23 \pmod{24}.\end{cases}$$

Proof. Taking $m=3$ and $t=\frac12$ in Theorem 2.1 we see that

$$\sum_{k=0}^{[p/6]}\binom{[\frac{p}{3}]}{k}\binom{[\frac{p}{6}]}{k}\frac{1}{2^k}\equiv\frac{1}{2^{[p/6]}}\sum_{k=0}^{[p/6]}\binom{[\frac{p}{3}]}{2k}\binom{\frac{p-1}{2}}{k}$$
$$\equiv\begin{cases}P_{\frac{p-1}{3}}(\frac{1}{\sqrt2}) \pmod p & \text{if } p\equiv 1 \pmod 3,\\ \sqrt2 P_{\frac{p-2}{3}}(\frac{1}{\sqrt2}) \pmod p & \text{if } p\equiv 2 \pmod 3.\end{cases}$$

From [S6, Theorem 4.5] we have

$$P_{[\frac{p}{3}]}\Big(\frac{1}{\sqrt2}\Big)\equiv\begin{cases}2x(\frac{x}{3}) \pmod p & \text{if } p=x^2+6y^2\equiv 1,7 \pmod{24},\\ 0 \pmod p & \text{if } p\equiv 17,23 \pmod{24}\end{cases}$$

Observe that $2^{-\frac{p-1}{6}}=2^{\frac{p-1}{3}-\frac{p-1}{2}}\equiv 2^{\frac{p-1}{3}} \pmod p$ for $p\equiv 1,7 \pmod{24}$. From the above we deduce the result.

Conjecture 4.2. *Let $p>3$ be a prime. Then*

$$\sum_{k=0}^{p-1}\frac{\binom{-\frac13}{k}\binom{-\frac16}{k}}{2^k}\equiv\Big(\frac{2}{p}\Big)2^{\frac{(\frac{p}{3})-1}{2}}\sum_{k=0}^{p-1}\frac{\binom{-\frac23}{k}\binom{-\frac56}{k}}{2^k}$$
$$\equiv\begin{cases}(\frac{x}{3})(2x-\frac{p}{2x}) \pmod {p^2} & \text{if } p=x^2+6y^2\equiv 1,7 \pmod{24},\\ (\frac{x}{3})(2x-\frac{p}{4x}) \pmod {p^2} & \text{if } p=2x^2+3y^2\equiv 5,11 \pmod{24},\\ 0 \pmod p & \text{if } p\equiv 13,19 \pmod{24}\end{cases}$$

and so

$$x\Big(\frac{x}{3}\Big)\equiv-\frac14\sum_{k=0}^{(p-5)/6}\binom{\frac{p-2}{3}}{k}\binom{\frac{p-5}{6}}{k}\frac{1}{2^k} \pmod p \quad \text{for} \quad p=2x^2+3y^2.$$

Theorem 4.3. *Let p be an odd prime such that $(\frac{17}{p})=1$. Then*

$$\sum_{k=0}^{[p/6]}\binom{[\frac p3]}{k}\binom{[\frac p6]}{k}\frac{1}{(-16)^k}$$
$$\equiv\begin{cases} x \pmod p & \text{if } p\equiv 1 \pmod 3 \text{ and so } 4p=x^2+51y^2 \text{ with } 3\mid x-2,\\ 0 \pmod p & \text{if } p\equiv 2 \pmod 3).\end{cases}$$

Proof. Taking $m=3$ and $t=\frac{17}{16}$ in Theorem 2.1 we see that

$$\sum_{k=0}^{[p/6]}\binom{[\frac p3]}{k}\binom{[\frac p6]}{k}\frac{1}{(-16)^k}\equiv\begin{cases} P_{[\frac p3]}(\frac{\sqrt{17}}{4}) \pmod p & \text{if } p\equiv 1 \pmod 3),\\ \frac{4}{\sqrt{17}}P_{[\frac p3]}(\frac{\sqrt{17}}{4}) \pmod p & \text{if } p\equiv 2 \pmod 3).\end{cases}$$

On the other hand, by [S6, Theorem 4.8],

$$P_{[\frac p3]}\Big(\frac{\sqrt{17}}{4}\Big)\equiv\begin{cases} -(\frac x3)x \pmod p & \text{if } p\equiv 1 \pmod 3) \text{ and so } 4p=x^2+51y^2,\\ 0 \pmod p & \text{if } p\equiv 2 \pmod 3).\end{cases}$$

Thus the theorem is proved.

Conjecture 4.3. *Let $p>3$ be a prime. Then*

$$\sum_{k=0}^{p-1}\frac{\binom{-\frac13}{k}\binom{-\frac16}{k}}{(-16)^k}\equiv\Big(\frac{17}{p}\Big)\Big(\frac{17}{16}\Big)^{\frac{1-(\frac p3)}{2}}\sum_{k=0}^{p-1}\frac{\binom{-\frac23}{k}\binom{-\frac56}{k}}{(-16)^k}$$
$$\equiv\begin{cases} -(\frac x3)(x-\frac px) \pmod{p^2} & \text{if } 4p=x^2+51y^2,\\ \frac14(\frac x3)(17x-\frac px) \pmod{p^2} & \text{if } 4p=17x^2+3y^2,\\ 0 \pmod p & \text{if } (\frac p3)=-(\frac{p}{17})=1\end{cases}$$

and so

$$x\Big(\frac x3\Big)\equiv-\frac14\sum_{k=0}^{(p-5)/6}\binom{\frac{p-2}{3}}{k}\binom{\frac{p-5}{6}}{k}\frac{1}{(-16)^k} \pmod p \quad \text{for} \quad 4p=17x^2+3y^2.$$

Using the theorems in Section 4 in [S6] and Theorem 2.1 one can similarly deduce the following results.

Theorem 4.4. *Let p be an odd prime such that $(\frac{41}{p})=1$. Then*

$$\sum_{k=0}^{[p/6]}\binom{[\frac p3]}{k}\binom{[\frac p6]}{k}\frac{1}{(-1024)^k}$$
$$\equiv\begin{cases} x \pmod p & \text{if } p\equiv 1 \pmod 3) \text{ and so } 4p=x^2+123y^2 \text{ with } 3\mid x-2,\\ 0 \pmod p & \text{if } p\equiv 2 \pmod 3).\end{cases}$$

Conjecture 4.4. *Let $p>3$ be a prime. Then*

$$\sum_{k=0}^{p-1}\frac{\binom{-\frac13}{k}\binom{-\frac16}{k}}{(-1024)^k}\equiv\Big(\frac{41}{p}\Big)\Big(\frac{1025}{1024}\Big)^{\frac{1-(\frac p3)}{2}}\sum_{k=0}^{p-1}\frac{\binom{-\frac23}{k}\binom{-\frac56}{k}}{(-1024)^k}$$

$$\equiv\begin{cases} -(\frac x3)(x-\frac px)\pmod{p^2} & \textit{if } 4p=x^2+123y^2,\\ \frac{5}{32}(\frac x3)(41x-\frac px)\pmod{p^2} & \textit{if } 4p=41x^2+3y^2,\\ 0\pmod p & \textit{if } (\frac p3)=-(\frac p{41})=1\end{cases}$$

and so

$$x\Big(\frac x3\Big)\equiv-\frac{5}{32}\sum_{k=0}^{(p-5)/6}\binom{\frac{p-2}{3}}{k}\binom{\frac{p-5}{6}}{k}\frac{1}{(-1024)^k}\pmod p\quad \textit{for}\quad 4p=41x^2+3y^2.$$

Theorem 4.5. *Let p be an odd prime such that $(\frac{89}{p})=1$. Then*

$$\sum_{k=0}^{[p/6]}\binom{[\frac p3]}{k}\binom{[\frac p6]}{k}\frac{1}{(-250000)^k}$$

$$\equiv\begin{cases} x\pmod p & \textit{if } p\equiv 1\pmod 3 \textit{ and so } 4p=x^2+267y^2 \textit{ with } 3\mid x-2,\\ 0\pmod p & \textit{if } p\equiv 2\pmod 3.\end{cases}$$

Conjecture 4.5. *Let $p>3$ be a prime. Then*

$$\sum_{k=0}^{p-1}\frac{\binom{-\frac13}{k}\binom{-\frac16}{k}}{(-250000)^k}\equiv\Big(\frac{89}{p}\Big)\Big(\frac{250001}{250000}\Big)^{\frac{1-(\frac p3)}{2}}\sum_{k=0}^{p-1}\frac{\binom{-\frac23}{k}\binom{-\frac56}{k}}{(-250000)^k}$$

$$\equiv\begin{cases} -(\frac x3)(x-\frac px)\pmod{p^2} & \textit{if } 4p=x^2+267y^2,\\ \frac{53}{500}(\frac x3)(89x-\frac px)\pmod{p^2} & \textit{if } 4p=89x^2+3y^2,\\ 0\pmod p & \textit{if } (\frac p3)=-(\frac p{89})=1\end{cases}$$

and so

$$x\Big(\frac x3\Big)\equiv-\frac{53}{500}\sum_{k=0}^{(p-5)/6}\binom{\frac{p-2}{3}}{k}\binom{\frac{p-5}{6}}{k}\frac{1}{(-250000)^k}\pmod p \text{ for } 4p=89x^2+3y^2.$$

Theorem 4.6. *Let p be a prime such that $p\equiv 1,4\pmod 5$. Then*

$$\sum_{k=0}^{[p/6]}\binom{[\frac p3]}{k}\binom{[\frac p6]}{k}\frac{1}{(-80)^k}$$

$$\equiv\begin{cases} x\pmod p & \textit{if } p\equiv 1,4\pmod{15} \textit{ and so } 4p=x^2+75y^2 \textit{ with } 3\mid x-2,\\ 0\pmod p & \textit{if } p\equiv 11,14\pmod{15}\end{cases}$$

Conjecture 4.6. *Let $p > 5$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac13}{k}\binom{-\frac16}{k}\frac{1}{(-80)^k}\\ &\equiv\Big(\frac{5}{p}\Big)\sum_{k=0}^{p-1}\binom{-\frac23}{k}\binom{-\frac56}{k}\frac{1}{(-80)^k}\\ &\equiv\begin{cases} x-\frac{p}{x} \pmod{p^2} & \text{if } p\equiv 1,19 \pmod{30} \text{ and so } 4p=x^2+75y^2 \text{ with } 3\mid x-2,\\ 5x-\frac{p}{5x} \pmod{p^2} & \text{if } p\equiv 7,13 \pmod{30} \text{ and so } 4p=25x^2+3y^2 \text{ with } 3\mid x-1,\\ 0 \pmod p & \text{if } p\equiv 17,23 \pmod{30}\end{cases}\end{aligned}$$

Theorem 4.7. *Let p be an odd prime and $p \neq 11$. Then*

$$\begin{aligned}&\sum_{k=0}^{[p/6]}\binom{[\frac p3]}{k}\binom{[\frac p6]}{k}\Big(\frac{27}{16}\Big)^k\\ &\equiv\begin{cases} -(\frac{-11+x/y}{p})(\frac{x}{11})x \pmod p & \text{if } (\frac p3)=(\frac p{11})=1 \text{ and so } 4p=x^2+11y^2,\\ 4y(\frac{-11+(\frac{x}{11})x/y}{p}) \pmod p & \text{if } (\frac p{11})=-(\frac p3)=1 \text{ and so } 4p=x^2+11y^2,\\ 0 \pmod p & \text{if } (\frac p{11})=-1.\end{cases}\end{aligned}$$

Conjecture 4.7. *Let $p > 11$ be a prime such that $(\frac{p}{11}) = 1$ and so $4p = x^2 + 11y^2$. Then*

$$\begin{aligned}\sum_{k=0}^{p-1}\binom{-\frac13}{k}\binom{-\frac16}{k}\Big(\frac{27}{16}\Big)^k&\equiv\Big(-\frac{11}{16}\Big)^{\frac{1-(\frac p3)}{2}}\sum_{k=0}^{p-1}\binom{-\frac23}{k}\binom{-\frac56}{k}\Big(\frac{27}{16}\Big)^k\\ &\equiv\begin{cases} -(\frac{-11+x/y}{p})(\frac{x}{11})(x-\frac{p}{x}) \pmod{p^2} & \text{if } 3\mid p-1,\\ -\frac14(\frac{-11+(\frac{x}{11})x/y}{p})(11y-\frac{p}{y}) \pmod{p^2} & \text{if } 3\mid p-2\end{cases}\end{aligned}$$

and so

$$\begin{aligned}&y\Big(\frac{-11+(\frac{x}{11})x/y}{p}\Big)\\ &\equiv\frac14\sum_{k=0}^{(p-5)/6}\binom{\frac{p-2}{3}}{k}\binom{\frac{p-5}{6}}{k}\Big(\frac{27}{16}\Big)^k \pmod p \quad \text{for} \quad 4p=x^2+11y^2\equiv 2 \pmod 3.\end{aligned}$$

Theorem 4.8. *Let $p>3$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{[p/6]}\binom{[\frac p3]}{k}\binom{[\frac p6]}{k}\Big(-\frac{9}{16}\Big)^k\\ &\equiv\begin{cases}L\pmod p & \text{if } p\equiv 1\pmod 3 \text{ and } 4p=L^2+27M^2 \text{ with } 3\mid L-2,\\ 0\pmod p & \text{if } p\equiv 2\pmod 3.\end{cases}\end{aligned}$$

Proof. From [S6, Theorem 3.2] we know that

$$P_{[\frac p3]}\Big(\frac54\Big)\equiv\begin{cases}L\pmod p & \text{if } p\equiv 1\pmod 3 \text{ and } 4p=L^2+27M^2 \text{ with } 3\mid L-2,\\ 0\pmod p & \text{if } p\equiv 2\pmod 3.\end{cases}$$

Thus taking $m=3$ and $t=\frac{25}{16}$ in Theorem 2.1 and then applying the above we deduce the result.

Conjecture 4.8. *Let $p\equiv 1\pmod 3$ be a prime and so $4p=L^2+27M^2$ with $3\mid L-2$. Then*

$$\sum_{k=0}^{p-1}\binom{-\frac13}{k}\binom{-\frac16}{k}\Big(-\frac{9}{16}\Big)^k\equiv\sum_{k=0}^{p-1}\binom{-\frac23}{k}\binom{-\frac56}{k}\Big(-\frac{9}{16}\Big)^k\equiv L-\frac{p}{L}\pmod{p^2}.$$

Theorem 4.9. *Let p be an odd prime. Then*

$$\begin{aligned}&\sum_{k=0}^{[p/6]}\binom{[\frac p3]}{k}\binom{[\frac p6]}{k}\Big(\frac{27}{2}\Big)^k\\ &\equiv\begin{cases}(-1)^{[\frac p8]}(\frac{-2-c/d}{p})2c\pmod p & \text{if } p=c^2+2d^2\equiv 1,19\pmod{24} \text{ and } 4\mid c-1,\\ -\frac45(-1)^{[\frac p8]}(\frac{2+c/d}{p})d\pmod p & \text{if } p=c^2+2d^2\equiv 11,17\pmod{24} \text{ and } 4\mid c-1,\\ 0\pmod p & \text{if } p\equiv 5,7\pmod 8.\end{cases}\end{aligned}$$

Proof. By [S6, Theorem 4.3],

$$\begin{aligned}&P_{[\frac p3]}(5/\sqrt{-2})\\ &\equiv\begin{cases}(-1)^{[\frac p8]}(\frac{-2-\sqrt{-2}}{p})2c\pmod p & \text{if } p=c^2+2d^2\equiv 1,3\pmod 8 \text{ and } 4\mid c-1,\\ 0\pmod p & \text{if } p\equiv 5,7\pmod 8)\end{cases}\end{aligned}$$

Thus taking $m=3$ and $t=-\frac{25}{2}$ in Theorem 2.1 and then applying the above we deduce the result.

Conjecture 4.9. *Let $p\equiv 1,3 \pmod 8$ be a prime and so $p=c^2+2d^2$ with $4\mid c-1$. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac13}{k}\binom{-\frac16}{k}\Big(\frac{27}{2}\Big)^k\\&\equiv\Big(-\frac{25}{2}\Big)^{\frac{1-(\frac p3)}{2}}\sum_{k=0}^{p-1}\binom{-\frac23}{k}\binom{-\frac56}{k}\Big(\frac{27}{2}\Big)^k\\&\equiv\begin{cases}(-1)^{[\frac p8]}(\frac{-2-c/d}{p})(2c-\frac{p}{2c}) \pmod {p^2} & \text{if } p\equiv 1,19 \pmod{24},\\ (-1)^{[\frac p8]}(\frac{2+c/d}{p})(10d-\frac{5p}{4d}) \pmod {p^2} & \text{if } p\equiv 11,17 \pmod{24}.\end{cases}\end{aligned}$$

5. Congruences for $\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}{8}]}{k}(1-t)^k \pmod p$.

Theorem 5.1. *Let $p>3$ be a prime and $t\in\mathbb{Z}_p$ with $t\not\equiv 0 \pmod p$. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac18}{k}\binom{-\frac38}{k}(1-t)^k\\&\equiv t^{\langle -\frac18\rangle_p}\sum_{k=0}^{p-1}\binom{-\frac18}{k}\binom{-\frac58}{k}\Big(1-\frac1t\Big)^k\\&\equiv\begin{cases}P_{[\frac p4]}(\sqrt t) \pmod p & \text{if } p\equiv 1,3 \pmod 8,\\ (\frac tp)\sqrt t P_{[\frac p4]}(\sqrt t) \pmod p & \text{if } p\equiv 5,7 \pmod 8.\end{cases}\end{aligned}$$

Proof. Taking $a=-\frac18$ in Theorem 2.2 and applying Lemmas 2.2 and 2.3 we see that

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac18}{k}\binom{-\frac38}{k}(1-t)^k\\&\equiv t^{\langle -\frac18\rangle_p}\sum_{k=0}^{p-1}\binom{-\frac18}{k}\binom{-\frac58}{k}\Big(1-\frac1t\Big)^k\equiv P_{2\langle -\frac18\rangle_p}(\sqrt t)\\&=\begin{cases}P_{\frac{p-1}{4}}(\sqrt t) \pmod p & \text{if } p\equiv 1 \pmod 8,\\ P_{\frac{3p-1}{4}}(\sqrt t)\equiv P_{\frac{p-3}{4}}(\sqrt t) \pmod p & \text{if } p\equiv 3 \pmod 8,\\ P_{\frac{5p-1}{4}}(\sqrt t)\equiv(\sqrt t)^pP_{\frac{p-1}{4}}(\sqrt t)\equiv(\frac tp)\sqrt tP_{\frac{p-1}{4}}(\sqrt t) \pmod p & \text{if } p\equiv 5 \pmod 8,\\ P_{\frac{7p-1}{4}}(\sqrt t)\equiv(\sqrt t)^pP_{\frac{3p-1}{4}}(\sqrt t)\equiv(\frac tp)\sqrt tP_{\frac{p-3}{4}}(\sqrt t) \pmod p & \text{if } p\equiv 7 \pmod 8.\end{cases}\end{aligned}$$

Thus the theorem is proved.

Theorem 5.2. *Let p be a prime such that $p\equiv \pm 1\pmod 8$. Then*

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\frac 1{9^k}$$
$$\equiv\begin{cases} 2x(\frac x3)\cdot 2^{\frac{p-1}4}\pmod p & \text{if } p=x^2+6y^2\equiv 1\pmod{24},\\ 3x(\frac x3)\cdot 2^{\frac{p-3}4}\pmod p & \text{if } p=x^2+6y^2\equiv 7\pmod{24},\\ 0\pmod p & \text{if } p\equiv 17,23\pmod{24}.\end{cases}$$

Proof. From [S7, Theorem 3.5] we know that

$$P_{[\frac p4]}\Big(\frac{2\sqrt 2}3\Big)\equiv\begin{cases}(-1)^{\frac{p-1}2}(\frac{\sqrt 2}p)(\frac x3)2x\pmod p & \text{if } p=x^2+6y^2\equiv 1,7\pmod{24},\\ 0\pmod p & \text{if } p\equiv 17,23\pmod{24}\end{cases}$$

Taking $m=4$ and $t=\frac 89$ in Theorem 2.1 and then applying the above we deduce

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\frac 1{9^k}$$
$$\equiv\begin{cases} P_{[\frac p4]}(\frac{2\sqrt 2}3)\equiv 2x(\frac x3)(\frac{\sqrt 2}p)\equiv 2x(\frac x3)\cdot 2^{\frac{p-1}4}\pmod p\\ \qquad\qquad\qquad \text{if } p=x^2+6y^2\equiv 1\pmod{24},\\ \frac 3{2\sqrt 2}P_{[\frac p4]}(\frac{2\sqrt 2}3)\equiv\frac 3{2\sqrt 2}(\frac{\sqrt 2}p)(\frac x3)2x\equiv 3x(\frac x3)\cdot 2^{\frac{p-3}4}\pmod p\\ \qquad\qquad\qquad \text{if } p=x^2+6y^2\equiv 7\pmod{24},\\ 0\pmod p\qquad \text{if } p\equiv 17,23\pmod{24}.\end{cases}$$

This proves the theorem.

Conjecture 5.1. *Let p be a prime such that $p\equiv 5,11\pmod{24}$ and so $p=2x^2+3y^2(x,y\in\mathbb Z)$. Then*

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\frac 1{9^k}\equiv\begin{cases}3x\frac ab\pmod p & \text{if } p=a^2+b^2\equiv 5\pmod{24},\\ 2x\frac cd\pmod p & \text{if } p=c^2+2d^2\equiv 11\pmod{24}.\end{cases}$$

Lemma 5.1 ([S4, Lemma 4.1]). *Let p be an odd prime. Then*

$$P_{[\frac p4]}(\sqrt t)\equiv-\sum_{n=0}^{p-1}(n^3+4n^2+2(1-\sqrt t)n)^{\frac{p-1}2}\pmod p.$$

Theorem 5.3. *Let* $p \equiv 1, 9 \pmod{20}$ *be a prime and so* $p = x^2 + 5y^2$ *with* $x, y \in \mathbb{Z}$. *Then*

$$\sum_{k=0}^{[p/8]} \binom{[\frac{p}{8}]}{k}\binom{[\frac{3p}{8}]}{k}\frac{1}{(-4)^k} \equiv \begin{cases} 2x \pmod{p} & \text{if } p \equiv 1, 9 \pmod{40}, \\ 4ya/b \pmod{p} & \text{if } p = a^2 + b^2 \equiv 21, 29 \pmod{40}. \end{cases}$$

Proof. From [LM, Theorem 11] we know that

$$\sum_{n=0}^{p-1} \Big(\frac{n^3 + 4n^2 + (2 - \sqrt{5})n}{p}\Big) = 2x.$$

Thus, taking $m = 4$ and $t = \frac{5}{4}$ in Theorem 2.1 and Lemma 5.1 we obtain

$$\begin{aligned} &\sum_{k=0}^{[p/8]} \binom{[\frac{p}{8}]}{k}\binom{[\frac{3p}{8}]}{k}\frac{1}{(-4)^k} \\ &\equiv \begin{cases} P_{[\frac{p}{4}]}(\sqrt{\frac{5}{4}}) \equiv 2x \pmod{p} & \text{if } p \equiv 1, 9 \pmod{40}, \\ \frac{2}{\sqrt{5}} P_{[\frac{p}{4}]}(\sqrt{\frac{5}{4}}) \equiv \frac{4}{5}\sqrt{5}x \equiv 4y\frac{a}{b} \pmod{p} & \text{if } p = a^2 + b^2 \equiv 21, 29 \pmod{40}. \end{cases} \end{aligned}$$

This is the result.

Theorem 5.4. *Let* p *be a prime such that* $p \equiv 1, 9, 11, 19 \pmod{40}$ *and so* $p = x^2 + 10y^2$ *with* $x, y \in \mathbb{Z}$. *Then*

$$2x \equiv \sum_{k=0}^{[p/8]} \binom{[\frac{p}{8}]}{k}\binom{[\frac{3p}{8}]}{k}\frac{1}{81^k} \pmod{p}.$$

Proof. From [LM] and Deuring's theorem we deduce that (see [S6])

$$\sum_{n=0}^{p-1} \Big(\frac{n^3 + 4n^2 + (2 - \frac{8}{9}\sqrt{5})n}{p}\Big) = 2x.$$

Thus, taking $m = 4$ and $t = \frac{80}{81}$ in Theorem 2.1 and then applying Lemma 5.1 and the above we deduce the result.

Theorem 5.5. *Suppose that* p *is a prime such that* $(\frac{-1}{p}) = (\frac{13}{p}) = 1$ *and so* $p = x^2 + 13y^2 (x, y \in \mathbb{Z})$. *Then*

$$\sum_{k=0}^{[p/8]} \binom{[\frac{p}{8}]}{k}\binom{[\frac{3p}{8}]}{k}\frac{1}{(-324)^k} \equiv \begin{cases} 2x \pmod{p} & \text{if } p \equiv 1 \pmod{8}, \\ \frac{36}{5}y\frac{a}{b} \pmod{p} & \text{if } p = a^2 + b^2 \equiv 5 \pmod{8}. \end{cases}$$

Proof. From [LM] and Deuring's theorem we deduce (see [S6])

$$\sum_{n=0}^{p-1}\Big(\frac{n^3+4n^2+2(1-\frac{5}{18}\sqrt{13})n}{p}\Big)=2x.$$

Now taking $m=4$ and $t=\frac{325}{324}$ in Theorem 2.1 and then applying Lemma 5.1 and the above we deduce

$$\begin{aligned}&\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\frac{1}{(-324)^k}\\ &\equiv\begin{cases}P_{\frac{p-1}{4}}(\frac{5\sqrt{13}}{18})\equiv 2x \pmod p\\ \qquad\quad \text{if } p\equiv 1 \pmod 8,\\ \frac{18}{5\sqrt{13}}P_{\frac{p-1}{4}}(\frac{5\sqrt{13}}{18})\equiv\frac{18}{5\sqrt{13}}\cdot 2x\equiv\frac{36x}{5\sqrt{-13}\sqrt{-1}}\equiv\frac{36}{5}y\frac ab \pmod p\\ \qquad\quad \text{if } p=a^2+b^2\equiv 5 \pmod 8.\end{cases}\end{aligned}$$

This completes the proof.

Remark 5.1 Let $d\in\{5,10,13\}$, and $f(d)=-4,81,-324$ according as $d=5,10,13$. Let p be a prime such that $p=x^2+dy^2\equiv 1 \pmod 8$. After reading the author's conjectures on

$$\sum_{k=0}^{p-1}\frac{\binom{-\frac13}{k}\binom{-\frac16}{k}}{m^k},\quad \sum_{k=0}^{p-1}\frac{\binom{-\frac14}{k}^2}{m^k},\quad \sum_{k=0}^{p-1}\frac{\binom{-\frac13}{k}^2}{m^k},\quad \sum_{k=0}^{p-1}\frac{\binom{-\frac12}{k}\binom{-\frac14}{k}}{m^k} \pmod{p^2},$$

the author's brother Z.W. Sun conjectured

$$\sum_{k=0}^{p-1}\binom{-\frac18}{k}\binom{-\frac38}{k}\frac{1}{f(d)^k}\equiv 2x-\frac{p}{2x} \pmod{p^2}.$$

Theorem 5.6. *Suppose that p is a prime such that $(\frac{-1}{p})=(\frac{37}{p})=1$ and so $p=x^2+37y^2 (x,y\in\mathbb{Z})$. Then*

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\frac{1}{(-882^2)^k}\equiv\begin{cases}2x \pmod p & \text{if } p\equiv 1 \pmod 8,\\ \frac{1764a}{145b}y \pmod p & \text{if } p=a^2+b^2\equiv 5 \pmod 8.\end{cases}$$

Proof. From [LM] and Deuring's theorem we deduce (see [S6])

$$\sum_{n=0}^{p-1}\Big(\frac{n^3+4n^2+2(1-\frac{145}{882}\sqrt{37})n}{p}\Big)=2x.$$

Now taking $m=4$ and $t=\frac{37\cdot 145^2}{882^2}$ in Theorem 2.1 and then applying Lemma 5.1 and the above we deduce

$$\begin{aligned}&\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\frac{1}{(-882^2)^k}\\ &\equiv\begin{cases}P_{\frac{p-1}{4}}(\frac{145\sqrt{37}}{882})\equiv 2x \pmod p\\ \qquad\qquad \text{if } p\equiv 1 \pmod 8,\\ \frac{882}{145\sqrt{37}}P_{\frac{p-1}{4}}(\frac{145\sqrt{37}}{882})\equiv \frac{882}{145\sqrt{37}}\cdot 2x\equiv \frac{1764x}{145bx/(ay)}=\frac{1764a}{145b}y \pmod p\\ \qquad\qquad \text{if } p=a^2+b^2\equiv 5 \pmod 8.\end{cases}\end{aligned}$$

This completes the proof.

Conjecture 5.2. *Suppose that p is a prime such that $(\frac{-1}{p})=(\frac{37}{p})=1$ and so $p=x^2+37y^2(x,y\in\mathbb{Z})$. Then*

$$\sum_{k=0}^{p-1}\binom{-\frac18}{k}\binom{-\frac38}{k}\frac{1}{(-882^2)^k}\equiv 2x-\frac{p}{2x} \pmod{p^2}.$$

Theorem 5.7. *Let p be a prime such that $(\frac{2}{p})=(\frac{-11}{p})=1$ and so $p=x^2+22y^2$ with $x,y\in\mathbb{Z}$. Then*

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\frac{1}{9801^k}\equiv\begin{cases}2x \pmod p & \text{if } p\equiv 1 \pmod 8,\\ \frac{99}{70}\cdot 2^{\frac{p+1}{4}}x \pmod p & \text{if } p\equiv 7 \pmod 8.\end{cases}$$

Proof. From [LM] and Deuring's theorem we deduce (see [S6])

$$\sum_{n=0}^{p-1}\Big(\frac{n^3+4n^2+2(1-\frac{70}{99}\sqrt2)n}{p}\Big)=2x.$$

Thus, taking $m=4$ and $t=\frac{9800}{9801}$ in Theorem 2.1 and then applying Lemma 5.1 and the above we deduce the result.

Conjecture 5.3. *Let p be a prime such that $(\frac{2}{p})=(\frac{-11}{p})=-1$ and so $p=2x^2+11y^2(x,y\in\mathbb{Z})$. Then*

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\frac{1}{9801^k}\equiv\begin{cases}2x\frac{c}{d} \pmod p & \text{if } p=c^2+2d^2\equiv 3 \pmod 8,\\ \frac{99}{35}x\frac{a}{b} \pmod p & \text{if } p=a^2+b^2\equiv 5 \pmod 8.\end{cases}$$

Theorem 5.8. *Let p be a prime such that $(\frac{-2}{p}) = (\frac{29}{p}) = 1$ and so $p = x^2 + 58y^2$ with $x, y \in \mathbb{Z}$. Then*

$$2x \equiv \sum_{k=0}^{[p/8]} \binom{[\frac{p}{8}]}{k} \binom{[\frac{3p}{8}]}{k} \frac{1}{99^{4k}} \pmod p.$$

Proof. From [LM] and Deuring's theorem we have (see [S6])

$$\sum_{n=0}^{p-1} \Big(\frac{n^3 + 4n^2 + 2(1 - \frac{1820}{99^2}\sqrt{29})n}{p}\Big) = 2x.$$

Thus, taking $m = 4$ and $t = \frac{29 \cdot 1820^2}{99^4}$ in Theorem 2.1 and then applying Lemma 5.1 and the above we deduce the result.

Conjecture 5.4. *Let p be a prime such that $(\frac{-2}{p}) = (\frac{29}{p}) = -1$ and so $p = 2x^2 + 29y^2 (x, y \in \mathbb{Z})$. Then*

$$y \equiv \frac{910}{9801} \sum_{k=0}^{[p/8]} \binom{[\frac{p}{8}]}{k} \binom{[\frac{3p}{8}]}{k} \frac{1}{99^{4k}} \pmod p.$$

Theorem 5.9. *Let p be a prime such that $p \equiv 1, 19 \pmod{24}$ and so $p = x^2 + 2y^2$. Then*

$$2x \equiv \sum_{k=0}^{[p/8]} \binom{[\frac{p}{8}]}{k} \binom{[\frac{3p}{8}]}{k} \frac{1}{2401^k} \pmod p.$$

Proof. From [S7, the proof of Theorem 3.9] we know that

$$\sum_{n=0}^{p-1} \Big(\frac{n^3 + 4n^2 + (2 - \frac{40}{49}\sqrt{6})n}{p}\Big) = 2x.$$

Thus, taking $m = 4$ and $t = \frac{2400}{2401}$ in Theorem 2.1 and then applying Lemma 5.1 and the above we deduce the result.

Conjecture 5.5. *let p be a prime such that $p \equiv 1, 3 \pmod 8$ and so $p = x^2 + 2y^2$. Then*

$$\sum_{k=0}^{p-1} \binom{-\frac{1}{8}}{k} \binom{-\frac{3}{8}}{k} \frac{1}{2401^k} \equiv 2x - \frac{p}{2x} \pmod{p^2}.$$

Theorem 5.10. *Let $p > 7$ be a prime. Then*

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\Big(\frac{256}{81}\Big)^k \equiv \begin{cases} 2x(\frac{3(7+x/y)}{p})(\frac x7) \pmod p & \text{if } p=x^2+7y^2\equiv 1,3 \pmod 8,\\ \frac{18}{5}y(\frac{3(7+(\frac x7)x/y)}{p}) \pmod p & \text{if } p=x^2+7y^2\equiv 5,7 \pmod 8,\\ 0 \pmod p & \text{if } p\equiv 3,5,6 \pmod 7.\end{cases}$$

Proof. From [S7, Theorem 3.1] we know that

$$P_{[\frac p4]}\Big(\frac{5\sqrt{-7}}{9}\Big)\equiv\begin{cases} 2x(\frac{3(7+x/y)}{p})(\frac x7) \pmod p & \text{if } p=x^2+7y^2\equiv 1,2,4 \pmod 7,\\ 0 \pmod p & \text{if } p\equiv 3,5,6 \pmod 7.\end{cases}$$

Now taking $m = 4$ and $t = -\frac{175}{81}$ in Theorem 2.1 and then applying the above we deduce the result.

Theorem 5.11. *Let $p > 7$ be a prime. Then*

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\Big(-\frac{256}{3969}\Big)^k \equiv \begin{cases} 2(\frac p3)(\frac x7)x \pmod p & \text{if } p=x^2+7y^2\equiv 1,3 \pmod 8,\\ -\frac{126}{65}(\frac p3)(\frac x7)x \pmod p & \text{if } p=x^2+7y^2\equiv 5,7 \pmod 8,\\ 0 \pmod p & \text{if } p\equiv 3,5,6 \pmod 7.\end{cases}$$

Proof. From [S4, Theorem 2.6] we know that

$$(5.1)\quad P_{[\frac p4]}\Big(\frac{65}{63}\Big)\equiv\begin{cases} (-1)^{[\frac p4]}2x(\frac p3)(\frac x7) \pmod p & \text{if } p=x^2+7y^2\equiv 1,2,4 \pmod 7,\\ 0 \pmod p & \text{if } p\equiv 3,5,6 \pmod 7.\end{cases}$$

Now taking $m = 4$ and $t = \frac{65^2}{63^2}$ in Theorem 2.1 and then applying (5.1) we deduce the result.

Theorem 5.12. *Let $p > 7$ be a prime. Then*

$$\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\Big(\frac{32}{81}\Big)^k \equiv \begin{cases} (\frac p3)2a \pmod p & \text{if } p=a^2+b^2\equiv 1 \pmod 8 \text{ and } 4\mid a-1,\\ \frac{18}{7}(\frac p3)a \pmod p & \text{if } p=a^2+b^2\equiv 5 \pmod 8 \text{ and } 4\mid a-1,\\ 0 \pmod p & \text{if } p\equiv 3 \pmod 4.\end{cases}$$

Proof. From [S4, Theorem 2.4] we know that

$$(5.2)\qquad P_{[\frac p4]}\Big(\frac79\Big)\equiv\begin{cases} (\frac p3)2a \pmod p & \text{if } p=a^2+b^2\equiv 1 \pmod 4 \text{ and } 4\mid a-1,\\ 0 \pmod p & \text{if } p\equiv 3 \pmod 4.\end{cases}$$

Now taking $m = 4$ and $t = \frac{49}{81}$ in Theorem 2.1 and then applying (5.2) we deduce the result.

Theorem 5.13. *Let $p>7$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{[p/8]}\binom{[\frac p8]}{k}\binom{[\frac{3p}8]}{k}\Big(-\frac{16}{9}\Big)^k\\&\equiv\begin{cases}2A\pmod p&\text{if } p=A^2+3B^2\equiv 1,19\pmod{24}\text{ and } 3\mid A-1,\\-\frac65A\pmod p&\text{if } p=A^2+3B^2\equiv 7,13\pmod{24}\text{ and } 3\mid A-1,\\0\pmod p&\text{if } p\equiv 2\pmod 3.\end{cases}\end{aligned}$$

Proof. From [S4, Theorem 2.5] we know that
(5.3)

$$P_{[\frac p4]}\Big(\frac53\Big)\equiv\begin{cases}(-1)^{[\frac p4]}2A\pmod p&\text{if } p=A^2+3B^2\equiv 1\pmod 3\text{ and } 3\mid A-1,\\0\pmod p&\text{if } p\equiv 2\pmod 3.\end{cases}$$

Now taking $m=4$ and $t=\frac{25}{9}$ in Theorem 2.1 and then applying (5.3) we deduce the result.

6. Congruences for $\sum_{k=0}^{p-1}\binom{-\frac13}{k}^2m^k$ and $\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2m^k\pmod p$.

Theorem 6.1. *Let $p>3$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac13}{k}^2 9^k\equiv\frac{1}{3^{[\frac p3]}}\sum_{k=0}^{p-1}\binom{-\frac13}{k}^2\frac{1}{9^k}\\&\equiv\begin{cases}L\pmod p&\text{if } p\equiv 1\pmod 3,\ 4p=L^2+27M^2\text{ and } 3\mid L-2,\\0\pmod p&\text{if } p\equiv 2\pmod 3.\end{cases}\end{aligned}$$

Proof. By Theorem 2.3 we have

$$\begin{aligned}&9^{\langle-\frac13\rangle_p}\sum_{k=0}^{p-1}\binom{-\frac13}{k}^2\frac{1}{9^k}\\&\equiv\sum_{k=0}^{p-1}\binom{-\frac13}{k}^2 9^k\equiv 8^{\langle-\frac13\rangle_p}P_{\langle-\frac13\rangle_p}\Big(\frac{10}{8}\Big)\\&=\begin{cases}8^{\frac{p-1}{3}}P_{\frac{p-1}{3}}(\frac54)\equiv P_{\frac{p-1}{3}}(\frac54)\pmod p&\text{if } p\equiv 1\pmod 3,\\8^{\frac{2p-1}{3}}P_{\frac{2p-1}{3}}(\frac54)\equiv 2P_{\frac{p-2}{3}}(\frac54)\pmod p&\text{if } p\equiv 2\pmod 3.\end{cases}\end{aligned}$$

From [S6, Theorem 3.2] we know that

$$P_{[\frac p3]}\Big(\frac54\Big)\equiv\begin{cases}L\pmod p&\text{if } p\equiv 1\pmod 3,\\0\pmod p&\text{if } p\equiv 2\pmod 3.\end{cases}$$

Since

$$9^{\langle-\frac13\rangle_p}=\begin{cases}9^{\frac{p-1}{3}}\equiv 3^{-\frac{p-1}{3}}\pmod p&\text{if } p\equiv 1\pmod 3,\\9^{\frac{2p-1}{3}}\equiv 3^{-\frac{p-2}{3}}\pmod p&\text{if } p\equiv 2\pmod 3,\end{cases}$$

combining all the above we deduce the result.

Conjecture 6.1. *Let p be a prime such that $p \equiv 1 \pmod 3$, $4p = L^2 + 27M^2 (L, M \in \mathbb{Z})$ and $L \equiv 2 \pmod 3$. Then*

$$\sum_{k=0}^{p-1} \binom{-\frac{1}{3}}{k}^2 9^k \equiv L - \frac{p}{L} \pmod{p^2}$$

Theorem 6.2. *Let p be an odd prime. Then*

$$\sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k}^2 (-8)^k$$
$$\equiv \begin{cases} (-1)^{\frac{p-1}{4}} 2x \pmod p & \text{if } p = x^2 + y^2 \equiv 1 \pmod 4 \text{ and } 4 \mid x - 1, \\ 0 \pmod p & \text{if } p \equiv 3 \pmod 4 \end{cases}$$

and

$$\sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k}^2 \frac{1}{(-8)^k}$$
$$\equiv \begin{cases} (-1)^{\frac{y}{4}} 2x \pmod p & \text{if } p = x^2 + y^2 \equiv 1 \pmod 8 \text{ and } 4 \mid x - 1, \\ (-1)^{\frac{y-2}{4}} 2y \pmod p & \text{if } p = x^2 + y^2 \equiv 5 \pmod 8 \text{ and } 2 \nmid x, \\ 0 \pmod p & \text{if } p \equiv 3 \pmod 4. \end{cases}$$

Proof. By Lemma 2.2 and Theorem 2.3 we have

$$\sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k}^2 (-8)^k$$
$$\equiv (-8)^{\langle -\frac{1}{4} \rangle_p} \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k}^2 \frac{1}{(-8)^k}$$
$$\equiv (-9)^{\langle -\frac{1}{4} \rangle_p} P_{\langle -\frac{1}{4} \rangle_p}\Big(\frac{7}{9}\Big) = 9^{\langle -\frac{1}{4} \rangle_p} P_{\langle -\frac{1}{4} \rangle_p}\Big(-\frac{7}{9}\Big)$$
$$= \begin{cases} 9^{\frac{p-1}{4}} P_{\frac{p-1}{4}}(-\frac{7}{9}) \equiv (\frac{3}{p}) P_{\frac{p-1}{4}}(-\frac{7}{9}) \pmod p & \text{if } p \equiv 1 \pmod 4, \\ 9^{\frac{3p-1}{4}} P_{\frac{3p-1}{4}}(-\frac{7}{9}) \equiv 9^{\frac{3p-1}{4}} P_{\frac{p-3}{4}}(-\frac{7}{9}) \pmod p & \text{if } p \equiv 3 \pmod 4. \end{cases}$$

From [S4, Theorem 2.4] we know that

$$P_{[\frac{p}{4}]}\Big(-\frac{7}{9}\Big) \equiv \begin{cases} (-1)^{\frac{p-1}{4}} (\frac{p}{3}) 2x \pmod p & \text{if } p = x^2 + y^2 \equiv 1 \pmod 4 \text{ and } 4 \mid x - 1, \\ 0 \pmod p & \text{if } p \equiv 3 \pmod 4. \end{cases}$$

When $p \equiv 1 \pmod 4$, we have $(-8)^{-\langle -\frac{1}{4} \rangle_p} = (-8)^{-\frac{p-1}{4}} \equiv (-2)^{\frac{p-1}{4}} \pmod p$. It is well known that (see [BEW])

$$2^{\frac{p-1}{4}} \equiv \begin{cases} (-1)^{\frac{y}{4}} \pmod p & \text{if } p \equiv 1 \pmod 8, \\ (-1)^{\frac{y-2}{4}} \frac{y}{x} \pmod p & \text{if } p \equiv 5 \pmod 8. \end{cases}$$

Now combining all the above we deduce the result.

Conjecture 6.2. *Let p be a prime of the form $4k+1$ and so $p=x^2+y^2$ with $4\mid x-1$. Then*

$$\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2(-8)^k\equiv(-1)^{\frac{p-1}{4}}\Big(2x-\frac{p}{2x}\Big)\pmod{p^2}$$

and

$$\sum_{k=0}^{p-1}\frac{\binom{-\frac14}{k}^2}{(-8)^k}\equiv\begin{cases}(-1)^{\frac y4}(2x-\frac{p}{2x})\pmod{p^2}&\text{if } p\equiv 1\pmod 8,\\(-1)^{\frac{y-2}{4}}(2y-\frac{p}{2y})\pmod{p^2}&\text{if } p\equiv 5\pmod 8.\end{cases}$$

Theorem 6.3. *Let $p>3$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2 4^k\equiv\frac{3-(-1)^{\frac{p-1}{2}}}{2}\Big(\frac 2p\Big)\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2\frac{1}{4^k}\\&\equiv\begin{cases}(-1)^{\frac{p-1}{4}+\frac{A-1}{2}}2A\pmod p&\text{if } p=A^2+3B^2\equiv 1\pmod{12},\\(-1)^{\frac{p+1}{4}}6B\pmod p&\text{if } p=A^2+3B^2\equiv 7\pmod{12}\text{ and } 4\mid B-1,\\0\pmod p&\text{if } p\equiv 2\pmod 3)\end{cases}\end{aligned}$$

Proof. By Lemma 2.2 and Theorem 2.3 we have

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2 4^k\\&\equiv 4^{\langle-\frac14\rangle_p}\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2\frac{1}{4^k}\\&\equiv 3^{\langle-\frac14\rangle_p}P_{\langle-\frac14\rangle_p}\Big(\frac53\Big)=(-3)^{\langle-\frac14\rangle_p}P_{\langle-\frac14\rangle_p}\Big(-\frac53\Big)\\&=\begin{cases}(-3)^{\frac{p-1}{4}}P_{\frac{p-1}{4}}(-\frac53)\pmod p&\text{if } p\equiv 1\pmod 4,\\(-3)^{\frac{3p-1}{4}}P_{\frac{3p-1}{4}}(-\frac53)\equiv(-3)^{-\frac{p-3}{4}}P_{\frac{p-3}{4}}(-\frac53)\pmod p&\text{if } p\equiv 3\pmod 4.\end{cases}\end{aligned}$$

From [S4, Theorem 2.5] we know that

$$P_{[\frac p4]}\Big(-\frac53\Big)\equiv\begin{cases}2A\pmod p&\text{if } p=A^2+3B^2\equiv 1\pmod 3\text{ and } 3\mid A-1,\\0\pmod p&\text{if } p\equiv 2\pmod 3.\end{cases}$$

Hence the result is true for $p\equiv 2\pmod 3$.

Now assume $p=A^2+3B^2\equiv 1\pmod 3$ and $A\equiv 1\pmod 3$. If $p\equiv 1\pmod{12}$, by [S2, p.1317] we have $3^{\frac{p-1}{4}}\equiv(-1)^{\frac{A-1}{2}}\pmod p$ and $4^{\langle-\frac14\rangle_p}=4^{\frac{p-1}{4}}\equiv(\frac 2p)=(-1)^{\frac{p-1}{4}}\pmod p$. Hence

$$\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2 4^k\equiv(-1)^{\frac{p-1}{4}}\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2\frac{1}{4^k}\equiv(-1)^{\frac{p-1}{4}+\frac{A-1}{2}}2A\pmod p.$$

If $p \equiv 7 \pmod{12}$ and $B \equiv 1 \pmod 4$, by [S2, p.1317] we have $3^{\frac{p-3}{4}} \equiv \frac{B}{A} \pmod p$. Since $4^{\langle -\frac14\rangle_p} = 4^{\frac{3p-1}{4}} = 2^{p-1+\frac{p+1}{2}} \equiv 2(\frac{2}{p}) \pmod p$, by the above we get

$$\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2 4^k \equiv 2\Big(\frac{2}{p}\Big)\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2\frac{1}{4^k} \equiv (-1)^{\frac{p-3}{4}}\frac{A}{B}\cdot 2A \equiv (-1)^{\frac{p+1}{4}}6B \pmod p.$$

Now combining all the above we deduce the result.

Conjecture 6.3. *Let* $p \equiv 1 \pmod 3$ *be a prime and so* $p = A^2 + 3B^2$. *Then*

$$\begin{aligned}\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2 4^k &\equiv \sum_{k=0}^{p-1}\binom{-\frac14}{k}\binom{-\frac12}{k}(-8)^k\\ &\equiv \begin{cases}(-1)^{\frac{p-1}{4}+\frac{A-1}{2}}(2A-\frac{p}{2A}) \pmod{p^2} & \text{if } p\equiv 1 \pmod{12},\\ (-1)^{\frac{p+1}{4}+\frac{B-1}{2}}(6B-\frac{p}{2B}) \pmod{p^2} & \text{if } p\equiv 7 \pmod{12}.\end{cases}\end{aligned}$$

and

$$\sum_{k=0}^{p-1}\frac{\binom{-\frac14}{k}^2}{4^k} \equiv \begin{cases}(-1)^{\frac{A-1}{2}}(2A-\frac{p}{2A}) \pmod{p^2} & \text{if } p\equiv 1 \pmod{12},\\ (-1)^{\frac{B-1}{2}}(3B-\frac{p}{4B}) \pmod{p^2} & \text{if } p\equiv 7 \pmod{12}.\end{cases}$$

Theorem 6.4. *Let* $p \neq 2, 7$ *be a prime. Then*

$$\begin{aligned}\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2 64^k &\equiv \frac{9-7(-1)^{\frac{p-1}{2}}}{2}\Big(\frac{2}{p}\Big)\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2\frac{1}{64^k}\\ &\equiv \begin{cases}(-1)^{\frac{p-1}{4}+\frac{x-1}{2}}2x \pmod p & \text{if } p = x^2+7y^2 \equiv 1 \pmod 4,\\ (-1)^{\frac{p+1}{4}+\frac{y-1}{2}}42y \pmod p & \text{if } p = x^2+7y^2 \equiv 3 \pmod 4,\\ 0 \pmod p & \text{if } p \equiv 3,5,6 \pmod 7.\end{cases}\end{aligned}$$

Proof. By Lemma 2.2 and Theorem 2.3 we have

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2 64^k\\ &\equiv 64^{\langle -\frac14\rangle_p}\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2\frac{1}{64^k}\\ &\equiv 63^{\langle -\frac14\rangle_p}P_{\langle -\frac14\rangle_p}\Big(\frac{65}{63}\Big) = (-63)^{\langle -\frac14\rangle_p}P_{\langle -\frac14\rangle_p}\Big(-\frac{65}{63}\Big)\\ &= \begin{cases}(-63)^{\frac{p-1}{4}}P_{\frac{p-1}{4}}(-\frac{65}{63}) \pmod p & \text{if } p\equiv 1 \pmod 4,\\ (-63)^{\frac{3p-1}{4}}P_{\frac{3p-1}{4}}(-\frac{65}{63}) \equiv (-63)^{-\frac{p-3}{4}}P_{\frac{p-3}{4}}(-\frac{65}{63}) \pmod p & \text{if } p\equiv 3 \pmod 4.\end{cases}\end{aligned}$$

By [S4, Theorem 2.6],

$$P_{[\frac{p}{4}]}\Big(-\frac{65}{63}\Big)\equiv\begin{cases}2x(\frac{p}{3})(\frac{x}{7})\pmod p & \text{if } p=x^2+7y^2\equiv 1,2,4\pmod 7,\\ 0\pmod p & \text{if } p\equiv 3,5,6\pmod 7.\end{cases}$$

Hence the result is true for $p\equiv 3,5,6\pmod 7$.

Now suppose $p\equiv 1,2,4\pmod 7$ and so $p=x^2+7y^2$ with $x,y\in\mathbb{Z}$. If $p\equiv 1\pmod 4$, by [S2, p.1317] we have $7^{\frac{p-1}{4}}\equiv(-1)^{\frac{x-1}{2}}(\frac{x}{7})\pmod p$ and so

$$\begin{aligned}&(-63)^{\frac{p-1}{4}}P_{\frac{p-1}{4}}(-\frac{65}{63})\\ &\equiv(-1)^{\frac{p-1}{4}}\Big(\frac{3}{p}\Big)\cdot(-1)^{\frac{x-1}{2}}\Big(\frac{x}{7}\Big)\cdot 2x\Big(\frac{p}{3}\Big)\Big(\frac{x}{7}\Big)=(-1)^{\frac{p-1}{4}+\frac{x-1}{2}}2x\pmod p.\end{aligned}$$

If $p\equiv 3\pmod 4$, by [S2, p.1317] we have $7^{\frac{p-3}{4}}\equiv(-1)^{\frac{y+1}{2}}(\frac{x}{7})\frac{y}{x}\pmod p$ and so

$$\begin{aligned}&(-63)^{-\frac{p-3}{4}}P_{\frac{p-3}{4}}(-\frac{65}{63})\\ &\equiv(-1)^{\frac{p-3}{4}}3\Big(\frac{3}{p}\Big)\cdot(-1)^{\frac{y+1}{2}}\Big(\frac{x}{7}\Big)\frac{x}{y}\cdot 2x\Big(\frac{p}{3}\Big)\Big(\frac{x}{7}\Big)\equiv(-1)^{\frac{p+1}{4}+\frac{y-1}{2}}42y\pmod p.\end{aligned}$$

Note that

$$64^{\langle-\frac14\rangle_p}=\begin{cases}64^{\frac{p-1}{4}}\equiv(\frac{2}{p})\pmod p & \text{if } p\equiv 1\pmod 4,\\ 64^{\frac{3p-1}{4}}\equiv 8(\frac{2}{p})\pmod p & \text{if } p\equiv 3\pmod 4.\end{cases}$$

Combining all the above we deduce the result.

Conjecture 6.4. *Let $p>2$ be a prime such that $p\equiv 1,2,4\pmod 7$ and so $p=x^2+7y^2$. Then*

$$\sum_{k=0}^{p-1}\binom{-\frac14}{k}^2 64^k\equiv\begin{cases}(\frac{2}{p})(-1)^{\frac{x-1}{2}}(2x-\frac{p}{2x})\pmod{p^2} & \text{if } p\equiv 1\pmod 4,\\ (\frac{2}{p})(-1)^{\frac{y-1}{2}}(42y-\frac{3p}{2y})\pmod{p^2} & \text{if } p\equiv 3\pmod 4\end{cases}$$

and

$$\sum_{k=0}^{p-1}\frac{\binom{-\frac14}{k}^2}{64^k}\equiv\begin{cases}(-1)^{\frac{x-1}{2}}(2x-\frac{p}{2x})\pmod{p^2} & \text{if } p\equiv 1\pmod 4,\\ \frac34(-1)^{\frac{y-1}{2}}(7y-\frac{p}{4y})\pmod{p^2} & \text{if } p\equiv 3\pmod 4.\end{cases}$$

Conjecture 6.5. *Let p be a prime such that $p\equiv 5,7\pmod 8$. Then*

$$\sum_{k=0}^{p-1}(-1)^k\binom{-\frac14}{k}^2\equiv\begin{cases}(-1)^{\frac{x+1}{2}}(2x-\frac{p}{2x})\pmod{p^2} & \text{if } p=x^2+2y^2\equiv 1\pmod 8,\\ (-1)^{\frac{y-1}{2}}(4y-\frac{p}{2y})\pmod{p^2} & \text{if } p=x^2+2y^2\equiv 3\pmod 8,\\ 0\pmod p & \text{if } p\equiv 5,7\pmod 8.\end{cases}$$

Conjecture 6.6. *Let p be an odd prime.*

(i) *If $p \equiv 1 \pmod 4$ and so $p = x^2 + y^2$ with $2 \nmid x$, then*

$$\sum_{k=0}^{p-1} \frac{\binom{-\frac14}{k}\binom{-\frac12}{k}}{4^k} \equiv \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \binom{-\frac12}{k}\binom{-\frac16}{k} 2^k$$
$$\equiv \begin{cases} (-1)^{\frac{p-1}{4}+\frac{x+1}{2}}(2x - \frac{p}{2x}) \pmod{p^2} & \text{if } 12 \mid p-1, \\ 2y - \frac{p}{2y} \pmod{p^2} & \text{if } 12 \mid p-5. \end{cases}$$

(ii) *If $p \equiv 3 \pmod 4$, then*

$$\sum_{k=0}^{p-1} \frac{\binom{-\frac14}{k}\binom{-\frac12}{k}}{4^k} \equiv 0 \pmod{p^2} \quad \text{and} \quad \sum_{k=0}^{p-1} \binom{-\frac12}{k}\binom{-\frac16}{k} 2^k \equiv 0 \pmod p.$$

Conjecture 6.7. *Let p be an odd prime. Then*

$$\sum_{k=0}^{p-1} \frac{\binom{-\frac14}{k}\binom{-\frac12}{k}}{(-3)^k} \equiv (-1)^{\frac{p-1}{4}} \sum_{k=0}^{p-1} \frac{\binom{-\frac14}{k}\binom{-\frac12}{k}}{81^k}$$
$$\equiv \begin{cases} 2x - \frac{p}{2x} \pmod{p^2} & \text{if } p = x^2 + y^2 \equiv 1 \pmod 4 \text{ and } 2 \nmid x, \\ 0 \pmod p & \text{if } p \equiv 3 \pmod 4. \end{cases}$$

Conjecture 6.8. *Let p be an odd prime. Then*

$$\sum_{k=0}^{p-1} \frac{\binom{-\frac14}{k}\binom{-\frac12}{k}}{(-80)^k}$$
$$\equiv \begin{cases} 2x - \frac{p}{2x} \pmod{p^2} & \text{if } p = x^2 + y^2 \equiv \pm 1 \pmod 5 \text{ and } 2 \nmid x, \\ 2y - \frac{p}{2y} \pmod{p^2} & \text{if } p = x^2 + y^2 \equiv \pm 2 \pmod 5 \text{ and } 2 \nmid x, \\ 0 \pmod p & \text{if } p \equiv 3 \pmod 4. \end{cases}$$

Conjecture 6.9. *Let $p > 5$ be a prime. Then*

$$\sum_{k=0}^{p-1} \binom{-\frac14}{k}\binom{-\frac12}{k} 2^k$$
$$= \begin{cases} 2x - \frac{p}{2x} \pmod{p^2} & \text{if } p = x^2 + 2y^2 \text{ with } x \equiv 1 \pmod 4, \\ 0 \pmod p & \text{if } p \equiv 5, 7 \pmod 8. \end{cases}$$

Conjecture 6.10. *Let $p>5$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{-\frac12}{k}\binom{-\frac13}{k}(-3)^k\equiv\sum_{k=0}^{p-1}\frac{\binom{-\frac12}{k}\binom{-\frac13}{k}}{(-27)^k}\equiv\sum_{k=0}^{p-1}\frac{\binom{-\frac12}{k}\binom{-\frac23}{k}}{(-4)^k}\\&\equiv\Big(\frac p5\Big)\sum_{k=0}^{p-1}\frac{\binom{-\frac12}{k}\binom{-\frac13}{k}}{5^k}\equiv\Big(\frac{-1}{p}\Big)\sum_{k=0}^{p-1}\binom{-\frac12}{k}\binom{-\frac13}{k}2^k\\&\equiv\begin{cases}2A-\frac{p}{2A}\pmod{p^2}&\text{if }p=A^2+3B^2\equiv1\pmod 3\text{ with }3\mid A-1,\\0\pmod p&\text{if }p\equiv2\pmod 3.\end{cases}\end{aligned}$$

Conjecture 6.11. *Let $p>5$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\frac{\binom{-\frac12}{k}\binom{-\frac13}{k}}{(-4)^k}\\&=\begin{cases}(\frac p5)(2A-\frac{p}{2A})\pmod{p^2}\\\quad\text{if }p=A^2+3B^2\equiv1\pmod 3\text{ with }5\mid AB\text{ and }3\mid A-1,\\(\frac p5)(A+3B-\frac{p}{A+3B})\pmod{p^2}\\\quad\text{if }p=A^2+3B^2\equiv1\pmod 3\text{ with }A/B\equiv-1,-2\pmod 5\text{ and }3\mid A-1,\\0\pmod p\qquad\text{if }p\equiv2\pmod 3.\end{cases}\end{aligned}$$

Conjecture 6.12. *Let $p>5$ be a prime. Then*

$$\sum_{k=0}^{p-1}\binom{-\frac12}{k}\binom{-\frac16}{k}\Big(-\frac{3}{125}\Big)^k\equiv\begin{cases}2A-\frac{p}{2A}\pmod{p^2}&\text{if }p=A^2+3B^2\equiv1\pmod 3,\\0\pmod p&\text{if }p\equiv2\pmod 3.\end{cases}$$

References

[AAR] G. Andrews, R. Askey, R. Roy, *Special Functions,*, Encyclopedia Math. Appl., vol. 71, Cambridge Univ. Press, Cambridge, 1999.

[B] H. Bateman, *Higher transcendental functions, Vol.II*, McGraw-Hill Book Co. Inc., 1953.

[BEW] B.C. Berndt, R.J. Evans and K.S. Williams, *Gauss and Jacobi Sums*, John Wiley & Sons, New York, 1998.

[BM] J. Brillhart and P. Morton, *Class numbers of quadratic fields, Hasse invariants of elliptic curves, and the supersingular polynomial*, J. Number Theory **106** (2004), 79-111.

[G] H.W. Gould, *Combinatorial Identities, A Standardized Set of Tables Listing 500 Binomial Coefficient Summations*, Morgantown, W. Va., 1972.

[I] N. Ishii, *Trace of Frobenius endomorphism of an elliptic curve with complex multiplication*, Bull. Austral. Math. Soc. **70** (2004), 125-142.

[JM] A. Joux et F. Morain, *Sur les sommes de caractères liées aux courbes elliptiques à multiplication complexe*, J. Number Theory **55** (1995), 108-128.

[LH] D. H. Lee and S. G. Hahn, *Gauss sums and binomial coefficients*, J. Number Theory **92** (2002), 257-271.

[LM] F. Leprévost and F. Morain, *Revêtements de courbes elliptiques à multiplication complexe par des courbes hyperelliptiques et sommes de caractères*, J. Number Theory **64** (1997), 165-182.

[M] P. Morton, *Explicit identities for invariants of elliptic curves*, J. Number Theory **120** (2006), 234-271.

[PV] R. Padma and S. Venkataraman, *Elliptic curves with complex multiplication and a character sum*, J. Number Theory **61** (1996), 274-282.

[PR] J.C. Parnami and A.R. Rajwade, *A new cubic character sum*, Acta Arith. **40** (1982), 347-356.

[R1] A.R. Rajwade, *The Diophantine equation* $y^2 = x(x^2 + 21Dx + 112D^2)$ *and the conjectures of Birch and Swinnerton-Dyer*, J. Austral. Math. Soc. Ser. A **24** (1977), 286-295.

[R2] A.R. Rajwade, *On a conjecture of Williams*, Bull. Soc. Math. Belg. Ser. B **36** (1984), 1-4.

[RPR] D.B. Rishi, J.C. Parnami and A.R. Rajwade, *Evaluation of a cubic character sum using the* $\sqrt{-19}$ *division points of the curve* $y^2 = x^3 - 2^3 \cdot 19x + 2 \cdot 19^2$, J. Number Theory **19** (1984), 184-194.

[S1] Z.H. Sun, *On the number of incongruent residues of* $x^4 + ax^2 + bx$ *modulo* p, J. Number Theory **119** (2006), 210-241.

[S2] Z.H. Sun, *On the quadratic character of quadratic units*, J. Number Theory **128** (2008), 1295-1335.

[S3] Z.H. Sun, *Congruences concerning Legendre polynomials*, Proc. Amer. Math. Soc. **139** (2011), 1915-1929.

[S4] Z.H. Sun, *Congruences concerning Legendre polynomials II*, arXiv:1012.3898. http://arxiv.org/abs/1012.3898.

[S5] Z.H. Sun, *Congruences concerning Legendre polynomials III*, arXiv:1012.4234. http://arxiv.org/abs/1012.4234.

[S6] Z.H. Sun, *Congruences involving* $\binom{2k}{k}^2\binom{3k}{k}m^{-k}$*,preprint*, arXiv:1104.2789v3. http://arxiv.org/abs/1104.2789.

[S7] Z.H. Sun, *Congruences involving* $\binom{2k}{k}^2\binom{4k}{2k}m^{-k}$*,preprint*, arXiv:1104.3047. http://arxiv.org/abs/1104.3047.

[W] K.S. Williams, *Evaluation of character sums connected with elliptic curves*, Proc. Amer. Math. Soc. **73** (1979), 291-299.